

References

- Agarwal, N., Pacher, P., Tegeder, I., Amaya, F., Constantin, C. E., Brenner, G. J., et al. (2007). Cannabinoids mediate analgesia largely via peripheral type 1 cannabinoid receptors in nociceptors. *Nature Neuroscience*, 10(7), 870–879.
- Amir, R., Kocsis, J. D., & Devor, M. (2005). Multiple interacting sites of ectopic spike electogenesis in primary sensory neurons. *Journal of Neuroscience*, 25(10), 2576–2585.
- Binshtok, A. M., Bean, B. P., & Woolf, C. J. (2007). Inhibition of nociceptors by TRPV1-mediated entry of impermeant sodium channel blockers. *Nature*, 449(7162), 607–610.
- Braz, J. M., & Basbaum, A. I. (2008). Genetically expressed transneuronal tracer reveals direct and indirect serotonergic descending control circuits. *Journal of Comparative Neurology*, 507(6), 1990–2003.
- Chatzigeorgiou, M., Yoo, S., Watson, J. D., Lee, W. H., Spencer, W. C., Kindt, K. S., et al. (2010). Specific roles for DEG/ENaC and TRP channels in touch and thermosensation in *C. elegans* nociceptors. *Nature Neuroscience*, 13(7), 861–868.
- Clark, A. K., Yip, P. K., Grist, J., Gentry, C., Staniland, A. A., Marchand, F., et al. (2007). Inhibition of spinal microglial cathepsin S for the reversal of neuropathic pain. *Proceedings of the National Academy of Sciences of the United States of America*, 104(25), 10655–10660.
- Costigan, M., Scholz, J., & Woolf, C. J. (2009). Neuropathic pain: A maladaptive response of the nervous system to damage. *Annual Review of Neuroscience*, 32, 1–32.
- Coull, J. A., Beggs, S., Boudreau, D., Boivin, D., Tsuda, M., Inoue, K., et al. (2005). BDNF from microglia causes the shift in neuronal anion gradient underlying neuropathic pain. *Nature*, 438(7070), 1017–1021.

- Dhaka, A., Viswanath, V., & Patapoutian, A. (2006). Trp ion channels and temperature sensation. *Annual Review of Neuroscience*, 29, 135–161.
- Diatchenko, L., Slade, G. D., Nackley, A. G., Bhalang, K., Sigurdsson, A., Belfer, I., et al. (2005). Genetic basis for individual variations in pain perception and the development of a chronic pain condition. *Human Molecular Genetics*, 14(1), 135–143.
- Dib-Hajj, S. D., Cummins, T. R., Black, J. A., & Waxman, S. G. (2010). Sodium channels in normal and pathological pain. *Annual Review of Neuroscience*, 33, 325–347.
- Dina, O. A., McCarter, G. C., de Coupade, C., & Levine, J. D. (2003). Role of the sensory neuron cytoskeleton in second messenger signaling for inflammatory pain. *Neuron*, 39(4), 613–624.
- Eippert, F., Finsterbusch, J., Bingel, U., & Buchel, C. (2009). Direct evidence for spinal cord involvement in placebo analgesia. *Science*, 326(5951), 404.
- Flor, H., Denke, C., Schaefer, M., & Grusser, S. (2001). Effect of sensory discrimination training on cortical reorganisation and phantom limb pain. *Lancet*, 357(9270), 1763–1764.
- Gardell, L. R., Vanderah, T. W., Gardell, S. E., Wang, R., Ossipov, M. H., Lai, J., et al. (2003). Enhanced evoked excitatory transmitter release in experimental neuropathy requires descending facilitation. *Journal of Neuroscience*, 23(23), 8370–8379.
- Hendrich, J., Van Minh, A. T., Hebllich, F., Nieto-Rostro, M., Watschinger, K., Striessnig, J., et al. (2008). Pharmacological disruption of calcium channel trafficking by the alpha2delta ligand gabapentin. *Proceedings of the National Academy of Sciences of the United States of America*, 105(9), 3628–3633.
- Honore, E. (2007). The neuronal background K₂P channels: Focus on TREK1. *Nature Reviews Neuroscience*, 8(4), 251–261.

- Ikeda, H., Stark, J., Fischer, H., Wagner, M., Drdla, R., Jager, T., et al. (2006). Synaptic amplifier of inflammatory pain in the spinal dorsal horn. *Science*, 312(5780), 1659–1662.
- Indo, Y., Tsuruta, M., Hayashida, Y., Karim, M. A., Ohta, K., Kawano, T., et al. (1996). Mutations in the TRKA/NGF receptor gene in patients with congenital insensitivity to pain with anhidrosis. *Nature Genetics*, 13(4), 485–488.
- Jiang, Y. Q., Xing, G. G., Wang, S. L., Tu, H. Y., Chi, Y. N., Li, J., et al. (2008). Axonal accumulation of hyperpolarization-activated cyclic nucleotide-gated cation channels contributes to mechanical allodynia after peripheral nerve injury in rat. *Pain*, 137(3), 495–506.
- Karashima, Y., Talavera, K., Everaerts, W., Janssens, A., Kwan, K. Y., Vennekens, R., et al. (2009). TRPA1 acts as a cold sensor *in vitro* and *in vivo*. *Proceedings of the National Academy of Sciences of the United States of America*, 106(4), 1273–1278.
- Kawasaki, Y., Xu, Z. Z., Wang, X., Park, J. Y., Zhuang, Z. Y., Tan, P. H., et al. (2008). Distinct roles of matrix metalloproteases in the early- and late-phase development of neuropathic pain. *Nature Medicine*, 14(3), 331–336.
- Kawasaki, Y., Zhang, L., Cheng, J. K., & Ji, R. R. (2008). Cytokine mechanisms of central sensitization: Distinct and overlapping role of interleukin-1beta, interleukin-6, and tumor necrosis factor-alpha in regulating synaptic and neuronal activity in the superficial spinal cord. *Journal of Neuroscience*, 28(20), 5189–5194.
- Kohama, I., Ishikawa, K., & Koocs, J. D. (2000). Synaptic reorganization in the substantia gelatinosa after peripheral nerve neuroma formation: Aberrant innervation of lamina II neurons by Abeta afferents. *Journal of Neuroscience*, 20(4), 1538–1549.

- Kwan, K. Y., Glazer, J. M., Corey, D. P., Rice, F. L., & Stucky, C. L. (2009). TRPA1 modulates mechanotransduction in cutaneous sensory neurons. *Journal of Neuroscience*, 29(15), 4808–4819.
- Lacroix-Fralish, M. L., & Mogil, J. S. (2009). Progress in genetic studies of pain and analgesia. *Annual Review of Pharmacology and Toxicology*, 49, 97–121.
- Latremoliere, A., & Woolf, C. J. (2009). Central sensitization: A generator of pain hypersensitivity by central neural plasticity. *Journal of Pain*, 10(9), 895–926.
- Mogil, J. S., Wilson, S. G., Chesler, E. J., Rankin, A. L., Nemmani, K. V., Lariviere, W. R., et al. (2003). The melanocortin-1 receptor gene mediates female-specific mechanisms of analgesia in mice and humans. *Proceedings of the National Academy of Sciences of the United States of America*, 100(8), 4867–4872.
- Olave, M. J., & Maxwell, D. J. (2003). Neurokinin-1 projection cells in the rat dorsal horn receive synaptic contacts from axons that possess alpha2C-adrenergic receptors. *Journal of Neuroscience*, 23(17), 6837–6846.
- Patapoutian, A., Tate, S., & Woolf, C. J. (2009). Transient receptor potential channels: Targeting pain at the source. *Nature Reviews Drug Discovery*, 8(1), 55–68.
- Pernia-Andrade, A. J., Kato, A., Witschi, R., Nyilas, R., Katona, I., Freund, T. F., et al. (2009). Spinal endocannabinoids and CB1 receptors mediate C-fiber-induced heterosynaptic pain sensitization. *Science*, 325(5941), 760–764.
- Samad, T. A., Moore, K. A., Sapirstein, A., Billet, S., Allchorne, A., Poole, S., et al. (2001). Interleukin-1beta-mediated induction of Cox-2 in the CNS contributes to inflammatory pain hypersensitivity. *Nature*, 410(6827), 471–475.
- Scholz, J., Broom, D. C., Youn, D. H., Mills, C. D., Kohno, T., Suter,

- M. R., et al. (2005). Blocking caspase activity prevents transsynaptic neuronal apoptosis and the loss of inhibition in lamina II of the dorsal horn after peripheral nerve injury. *Journal of Neuroscience*, 25(32), 7317–7323.
- Scholz, J., & Woolf, C. J. (2007). The neuropathic pain triad: Neurons, immune cells and glia. *Nature Neuroscience*, 10(11), 1361–1368.
- Stein, C., & Lang, L. J. (2009). Peripheral mechanisms of opioid analgesia. *Current Opinion in Pharmacology*, 9(1), 3–8.
- Suzuki, R., Morcuende, S., Webber, M., Hunt, S. P., & Dickenson, A. H. (2002). Superficial NK1-expressing neurons control spinal excitability through activation of descending pathways. *Nature Neuroscience*, 5(12), 1319–1326.
- Tao, Y. X., Rumbaugh, G., Wang, G. D., Petralia, R. S., Zhao, C., Kauer, F. W., et al. (2003). Impaired NMDA receptor-mediated postsynaptic function and blunted NMDA receptor-dependent persistent pain in mice lacking postsynaptic density-93 protein. *Journal of Neuroscience*, 23(17), 6703–6712.
- Tegeder, I., Costigan, M., Griffin, R. S., Abele, A., Belfer, I., Schmidt, H., et al. (2006). GTP cyclohydrolase and tetrahydrobiopterin regulate pain sensitivity and persistence. *Nature Medicine*, 12(11), 1269–1277.
- Wemmie, J. A., Price, M. P., & Welsh, M. J. (2006). Acid-sensing ion channels: Advances, questions and therapeutic opportunities. *Trends in Neurosciences*, 29(10), 578–586.
- Woolf, C. J., & Ma, Q. (2007). Nociceptors—noxious stimulus detectors. *Neuron*, 55(3), 353–364.
- Woolf, C. J., Shortland, P., & Coggeshall, R. E. (1992). Peripheral nerve injury triggers central sprouting of myelinated afferents. *Nature*, 355(6355), 75–78.

Yaksh, T. L. (2006). Calcium channels as therapeutic targets in neuropathic pain. *Journal of Pain*, 7(1 Suppl. 1), S13–30.