

References

- Armstrong, N., Sun, Y., Chen, G. Q., & Gouaux, E. (1998). Structure of a glutamate-receptor ligand-binding core in complex with kainate. *Nature*, *395*, 913–917.
- Attwell, D., & Laughlin, S. B. (2001). An energy budget for signaling in the grey matter of the brain. *Journal of Cerebral Blood Flow and Metabolism*, *21*, 1133–1145.
- Baker, D. A., McFarland, K., Lake, R. W., et al. (2003). Neuroadaptations in cystine–glutamate exchange underlie cocaine relapse. *Nature Neuroscience*, *6*, 743–749.
- Bennett, J. A., & Dingledine, R. (1995). Topology profile for a glutamate receptor: three transmembrane domains and a channel-lining reentrant membrane loop. *Neuron*, *14*, 373–384.
- Braitenberg, V., & Schüz, A. (1998). *Cortex: Statistics and Geometry of Neuronal Connectivity* (2nd ed.). Berlin: Springer.
- Broman, J., Hassel, B., Rinvik, E., & Ottersen, O. P. (2000). Biochemistry and anatomy of transmitter glutamate. In O. P. Ottersen & J. Storm-Mathisen (Eds.), *Handbook of Chemical Neuroanatomy 18* (pp. 1–44). Elsevier.
- Brusa, R., Zimmermann, F., Koh, D. S., et al. (1995). Early-onset epilepsy and postnatal lethality associated with an editing-deficient GluR-B allele in mice. *Science*, *270*, 1677–1680.
- Chaudhry, F. A., Schmitz, D., Reimer, R. J., et al. (2002). Glutamine uptake by neurons: interaction of protons with system a transporters. *Journal of Neuroscience*, *22*, 62–72.
- Collingridge, G. L., Peineau, S., Howland, J. G., & Wang, Y. T. (2010). Long-term depression in the CNS. *Nature Reviews Neuroscience*, *11*, 459–473.
- Conn, P. J. (2003). Physiological roles and therapeutic potential of metabotropic glutamate receptors. *Annals of the New York Academy of Sciences*, *1003*, 12–21.

- Correia, S. S., Bassani, S., Brown, T. C., et al. (2008). Motor protein-dependent transport of AMPA receptors into spines during long-term potentiation. *Nature Neuroscience*, *11*, 457–466.
- Danbolt, N. C. (2001). Glutamate uptake. *Progress in Neurobiology*, *65*, 1–105.
- Ehlers, M. D., Zhang, S., Bernhardt, J. P., & Huganir, R. L. (1996). Inactivation of NMDA receptors by direct interaction of calmodulin with the NR1 subunit. *Cell*, *84*, 745–755.
- Fletcher, E. J., & Lodge, D. (1996). New developments in the molecular pharmacology of α -amino-3-hydroxy-5-methyl-4-isoxazole propionate and kainate receptors. *Pharmacology & Therapeutics*, *70*, 65–89.
- Freneau, R. T., Jr., Voglmaier, S., Seal, R. P., & Edwards, R. H. (2004). VGLUTs define subsets of excitatory neurons and suggest novel roles for glutamate. *Trends in Neurosciences*, *27*, 98–103.
- Hardingham, G. E., & Bading, H. (2010). Synaptic versus extrasynaptic NMDA receptor signalling: implications for neurodegenerative disorders. *Nature Reviews Neuroscience*, *11*, 682–696.
- Hering, H., Lin, C. C., & Sheng, M. (2003). Lipid rafts in the maintenance of synapses, dendritic spines, and surface AMPA receptor stability. *Journal of Neuroscience*, *23*, 3262–3271.
- Higuchi, M., Single, F. N., Kohler, M., et al. (1993). RNA editing of AMPA receptor subunit GluRB: a base-paired intron-exon structure determines position and efficiency. *Cell*, *75*, 1361–1370.
- Hollmann, M., O'Shea-Greenfield, A., Rogers, S. W., & Heinemann, S. (1989). Cloning by functional expression of a member of the glutamate receptor family. *Nature*, *342*, 643–648.

- Hong, S. J., Li, H., Becker, K. G., Dawson, V. L., & Dawson, T. M. (2004). Identification and analysis of plasticity-induced late-response genes. *Proceedings of the National Academy of Sciences of the USA*, *101*, 2145–2150.
- Hume, R. I., Dingledine, R., & Heinemann, S. F. (1991). Identification of a site in glutamate receptor subunits that controls calcium permeability. *Science*, *253*, 1028–1032.
- Jia, Z., Agopyan, N., Miu, P., et al. (1996). Enhanced LTP in mice deficient in the AMPA receptor GluR2. *Neuron*, *17*, 945–956.
- Jin, S. X., & Feig, L. A. (2010). Long-term potentiation in the CA1 hippocampus induced by NR2A subunit-containing NMDA glutamate receptors is mediated by Ras-GRF2/Erk map kinase signaling. *PLoS One*, *5*, e11732.
- Kano, M., Hashimoto, K., Kurihara, H., et al. (1997). Persistent multiple climbing fiber innervation of cerebellar Purkinje cells in mice lacking mGluR1. *Neuron*, *18*, 71–79.
- Kato, A. S., Gill, M. B., Yu, H., Nisenbaum, E. S., & Brecht, D. S. (2010). TARPs differentially decorate AMPA receptors to specify neuropharmacology. *Trends in Neurosciences*, *33*, 241–248.
- Kennedy, M. B., Beale, H. C., Carlisle, H. J., & Washburn, L. R. (2005). Integration of biochemical signalling in spines. *Nature Reviews Neuroscience*, *6*, 423–434.
- Kleckner, N. W., & Dingledine, R. (1988). Requirement for glycine in activation of NMDA receptors expressed in *Xenopus* oocytes. *Science*, *241*, 835–837.
- Kuner, T., Wollmuth, L. P., Karlin, A., Seeburg, P. H., & Sakmann, B. (1996). Structure of the NMDA receptor channel M2 segment inferred from the accessibility of substituted cysteines. *Neuron*, *17*, 343–352.
- Lin, D. T., Makino, Y., Sharma, K., et al. (2009). Regulation of AMPA receptor extrasynaptic

- insertion by 4.1N, phosphorylation and palmitoylation. *Nature Neuroscience*, *12*, 879–887.
- Mayer, M. L., & Armstrong, N. (2004). Structure and function of glutamate receptor ion channels. *Annual Review of Physiology*, *66*, 161–181.
- Nakanishi, S. (1992). Molecular diversity of glutamate receptors and implications for brain function. [Review]. *Science*, *258*(5082), 597–603.
- Nowak, L., Bregestovski, P., Ascher, P., et al. (1984). Magnesium gates glutamate-activated channels in mouse central neurones. *Nature*, *307*, 462–465.
- Pekhletski, R., Gerlai, R., Overstreet, L. S., et al. (1996). Impaired cerebellar synaptic plasticity and motor performance in mice lacking the mGluR4 subtype of metabotropic glutamate receptor. *Journal of Neuroscience*, *16*, 6364–6373.
- Shen, J., Petersen, K. F., Behar, K. L., et al. (1999). Determination of the rate of the glutamate/glutamine cycle in the human brain by in vivo ¹³C NMR. *Proceedings of the National Academy of Sciences of the USA*, *96*, 8235–8240.
- Si, K., Giustetto, M., Etkin, A., et al. (2003). A neuronal isoform of CPEB regulates local protein synthesis and stabilizes synapse-specific long-term facilitation in aplysia. *Cell*, *115*, 893–904.
- Skrede, K. K., & Malthe-Sørensen, D. (1981). Increased resting and evoked release of transmitter following repetitive electrical tetanization in hippocampus: a biochemical correlate to long-lasting synaptic potentiation. *Brain Research*, *208*, 436–441.
- Sommer, B., Keinänen, K., Verdoorn, T. A., et al. (1990). Flip and flop: a cell specific functional switch in glutamate-operated channels of the CNS. *Science*, *249*, 1580–1585.
- Takahashi, T., Forsythe, I. D., Tsujimoto, T., Barnes-Davies, M., & Onodera, K. (1996).

- Presynaptic calcium current modulation by a metabotropic glutamate receptor. *Science*, 274, 594–597.
- Tong, G., Shepard, D., & Jahr, C. E. (1995). Synaptic desensitization of NMDA receptors by calcineurin. *Science*, 267, 1510–1512.
- Traynelis, S. F., Wollmuth, L., Menniti, F., McBain, C., & Dingledine, R. (2010). Glutamate receptor ion channels: structure, regulation and function. *Pharmacological Reviews*, 62, 405–496.
- Traynelis, S. F., Hartley, M., & Heinemann, S. F. (1995). Control of proton sensitivity of the NMDA receptor by RNA splicing and polyamines. *Science*, 268, 873–876.
- Tsien, J. Z., Huerta, P. T., & Tonegawa, S. (1996). The essential role of hippocampal CA1 NMDA receptor-dependent synaptic plasticity in spatial memory. *Cell*, 87, 1327–1338.
- Voutsinos-Porche, B., Bonvento, G., Tanaka, K., et al. (2003). Glial glutamate transporters mediate a functional metabolic crosstalk between neurons and astrocytes in the mouse developing cortex. *Neuron*, 37, 275–286.
- Wollmuth, L. P., Kuner, T., Seeburg, P. H., & Sakmann, B. (1996). Differential contribution of the NR1- and NR2A-subunits to the selectivity filter of recombinant NMDA receptor channels. *Journal of Physiology*, 491, 779–797.
- Wolosker, H., Blackshaw, S., & Snyder, S. H. (1999). Serine racemase: a glial enzyme synthesizing D-serine to regulate glutamate-*N*-methyl-D-aspartate neurotransmission. *Proceedings of the National Academy of Sciences of the USA*, 96, 13409–13414.
- Yokoi, M., Kobayashi, K., Manabe, T., et al. (1996). Impairment of hippocampal mossy fiber LTD in mice lacking mGluR2. *Science*, 273, 645–647.