

## References

- Ascherio, A., LeWitt, P. A., Xu, K., Eberly, S., Watts, A., Matson, W. R., et al. (2009). Urate as a predictor of the rate of clinical decline in Parkinson disease. *Archives of Neurology*, *66*, 1460–1468.
- Barkhoudarian, M. T. & Schwarzschild, M. A. (2011). Preclinical jockeying on the translational track of adenosine A<sub>(2A)</sub> receptors. *Experimental Neurology* (In press).
- Berne, R. M. (1963). Cardiac nucleotides in hypoxia: Possible role in regulation of coronary blood flow. *American Journal of Physiology*, *204*, 317–322.
- Burnstock, G. (1978). A basis for distinguishing two types of purinergic receptor. In L. Bolis & R. W. Straub (Eds.), *Cell Membrane Receptors for Drugs and Hormones: A Multidisciplinary Approach* (pp. 107–118). New York: Raven Press.
- Carta, A. R., Kachroo, A., Schintu, N., Xu, K., Schwarzschild, M. A., Wardas, J., et al. (2009). Inactivation of neuronal forebrain A receptors protects dopaminergic neurons in a mouse model of Parkinson's disease. *Journal of Neurochemistry*, *111*, 1478–1489.
- Chekeni, F. B., Elliott, M. R., Sandilos, J. K., Walk, S. F., Kinchen, J. M., Lazarowski, E. R., et al. (2010). Pannexin-1 channels mediate 'find-me' signal release and membrane permeability during apoptosis. *Nature*, *467*, 863–867.
- Choi, D. S., Cascini, M. G., Mailliard, W., Young, H., Paredes, P., McMahon, T., et al. (2004). The type 1 equilibrative nucleoside transporter regulates ethanol intoxication and preference. *Nature Neuroscience*, *7*, 855–861.
- Cotrina, M. L., Lin, J. H., Alves-Rodrigues, A., Liu, S., Li, J., Azmi-Ghadimi, H., et al. (1998). Connexins regulate calcium signaling by controlling ATP release. *Proceedings of the National Academy of Sciences of the United States of America*, *95*, 15735–15740.

- Coull, J. A., Beggs, S., Boudreau, D., Boivin, D., Tsuda, M., Inoue, K., et al. (2005). BDNF from microglia causes the shift in neuronal anion gradient underlying neuropathic pain. *Nature*, *438*, 1017–1021.
- Davalos, D., Grutzendler, J., Yang, G., Kim, J. V., Zuo, Y., Jung, S., et al. (2005). ATP mediates rapid microglial response to local brain injury *in vivo*. *Nature Neuroscience*, *8*, 752–758.
- Dolphin, A. C., Forda, S. R., & Scott, R. H. (1986). Calcium-dependent currents in cultured rat dorsal root ganglion neurones are inhibited by an adenosine analogue. *Journal of Physiology*, *373*, 47–61.
- Drury, A. N., & Szent-Gyorgyi, A. (1929). The physiological activity of adenine compounds with especial reference to their action upon the mammalian heart. *Journal of Physiology*, *68*, 213–237.
- Dunwiddie, T. V., Diao, L., & Proctor, W. R. (1997). Adenine nucleotides undergo rapid, quantitative conversion to adenosine in the extracellular space in rat hippocampus. *Journal of Neuroscience*, *17*, 7673–7682.
- Ferre, S., O'Connor, W. T., Snaprud, P., Ungerstedt, U., & Fuxe, K. (1994). Antagonistic interaction between adenosine A<sub>2A</sub> receptors and dopamine D2 receptors in the ventral striopallidal system. Implications for the treatment of schizophrenia. *Neuroscience*, *63*, 765–773.
- Fields, R. D., & Ni, Y. (2010). Nonsynaptic communication through ATP release from volume-activated anion channels in axons. *Science Signaling*, *3*, ra73.
- Gao, X., Chen, H., Choi, H. K., Curhan, G., Schwarzschild, M. A., & Ascherio, A. (2008). Diet, urate, and Parkinson's disease risk in men. *American Journal of Epidemiology*, *167*, 831–

838.

- Gordon, G. R., Baimoukhametova, D. V., Hewitt, S. A., Rajapaksha, W. R., Fisher, T. E., & Bains, J. S. (2005). Norepinephrine triggers release of glial ATP to increase postsynaptic efficacy. *Nature Neuroscience*, *8*, 1078–1086.
- Gourine, A. V., Kasymov, V., Marina, N., Tang, F., Figueiredo, M. F., Lane, S., et al. (2010). Astrocytes control breathing through pH-dependent release of ATP. *Science*, *329*, 571–575.
- Guthrie, P. B., Knappenberger, J., Segal, M., Bennett, M. V., Charles, A. C., & Kater, S. B. (1999). ATP released from astrocytes mediates glial calcium waves. *Journal of Neuroscience*, *19*, 520–528.
- Halassa, M. M., Florian, C., Fellin, T., Munoz, J. R., Lee, S. Y., Abel, T., et al. (2009). Astrocytic modulation of sleep homeostasis and cognitive consequences of sleep loss. *Neuron*, *61*, 213–219.
- Khakh, B. S., Burnstock, G., Kennedy, C., King, B. F., North, R. A., Seguela, P., et al. (2001). International union of pharmacology. XXIV. Current status of the nomenclature and properties of P2X receptors and their subunits. *Pharmacological Reviews*, *53*, 107–118.
- Koizumi, S., Shigemoto-Mogami, Y., Nasu-Tada, K., Shinozaki, Y., Ohsawa, K., Tsuda, M., et al. (2007). UDP acting at P2Y6 receptors is a mediator of microglial phagocytosis. *Nature*, *446*, 1091–1095.
- Linden, J. (1991). Structure and function of A1 adenosine receptors. *The FASEB Journal*, *5*, 2668–2676.
- Maenhaut, C., Van Sande, J., Libert, F., Abramowicz, M., Parmentier, M., Vanderhaegen, J. J., et al. (1990). RDC8 codes for an adenosine A2 receptor with physiological constitutive

- activity. *Biochemical and Biophysical Research Communications*, *173*, 1169–1178.
- Manzoni, O. J., Manabe, T., & Nicoll, R. A. (1994). Release of adenosine by activation of NMDA receptors in the hippocampus. *Science*, *265*, 2098–2101.
- Pascual, O., Casper, K. B., Kubera, C., Zhang, J., Revilla-Sanchez, R., Sul, J. Y., et al. (2005). Astrocytic purinergic signaling coordinates synaptic networks. *Science*, *310*, 113–116.
- Porkka-Heiskanen, T., Strecker, R. E., Thakkar, M., Bjorkum, A. A., Greene, R. W., & McCarley, R. W. (1997). Adenosine: A mediator of the sleep-inducing effects of prolonged wakefulness. *Science*, *276*, 1265–1268.
- Resta, R., & Thompson, L. F. (1997). T cell signalling through CD73. *Cellular Signalling*, *9*, 131–139.
- Richardson, P. J., Kase, H., & Jenner, P. G. (1997). Adenosine A<sub>2A</sub> receptor antagonists as new agents for the treatment of Parkinson's disease. *Trends in Pharmacological Sciences*, *18*, 338–344.
- Sawada, K., Echigo, N., Juge, N., Miyaji, T., Otsuka, M., Omote, H., et al. (2008). Identification of a vesicular nucleotide transporter. *Proceedings of the National Academy of Sciences of the United States of America*, *105*, 5683–5686.
- Serrano, A., Haddjeri, N., Lacaille, J. C., & Robitaille, R. (2006). GABAergic network activation of glial cells underlies hippocampal heterosynaptic depression. *Journal of Neuroscience*, *26*, 5370–5382.
- Stevens, B., & Fields, R. D. (2000). Response of Schwann cells to action potentials in development. *Science*, *287*, 2267–2271.
- Stevens, B., Porta, S., Haak, L. L., Gallo, V., & Fields, R. D. (2002). Adenosine: a neuron—glial transmitter promoting myelination in the CNS in response to action potentials. *Neuron*,

36, 855–868.

- Szybala, C., Pritchard, E. M., Lusardi, T. A., Li, T., Wilz, A., Kaplan, D. L., et al. (2009). Antiepileptic effects of silk-polymer based adenosine release in kindled rats. *Experimental Neurology*, *219*, 126–135.
- Trussell, L. O., & Jackson, M. B. (1985). Adenosine-activated potassium conductance in cultured striatal neurons. *Proceedings of the National Academy of Sciences of the United States of America*, *82*, 4857–4861.
- Wang, Z., Haydon, P. G., & Yeung, E. S. (2000). Direct observation of calcium-independent intercellular ATP signaling in astrocytes. *Analytical Chemistry*, *72*, 2001–2007.
- Zimmermann, H., & Braun, N. (1996). Extracellular metabolism of nucleotides in the nervous system. *Journal of Autonomic Pharmacology*, *16*, 397–400.