

## References

- Aloy, P., Companys, V., Vendrell, J., Aviles, F. X., Fricker, L. D., Coll, M., et al. (2001). The crystal structure of the inhibitor-complexed carboxypeptidase D domain II and the modeling of regulatory carboxypeptidases. *Journal of Biological Chemistry*, 276, 16177–16184.
- Arolas, J. L., Vendrell, J., Aviles, F. X., & Fricker, L. D. (2007). Metalloproteinases: Emerging drug targets in biomedicine. *Current Pharmaceutical Design*, 13, 347–364.
- Brothers, S. P., Janovick, J. A., & Conn, P. M. (2003). Unexpected effects of epitope and chimeric tags on gonadotropin-releasing hormone receptors: Implications for understanding the molecular etiology of hypogonadotropic hypogonadism. *Journal of Clinical Endocrinology and Metabolism*, 88, 6107–6112.
- Chufan, E. E., De, M., Eipper, B. A., Mains, R. E., & Amzel, L. M. (2010). Amidation of bioactive peptides: The structure of the lyase domain of the amidating enzyme. *Structure*, 17, 965–973.
- Czyzyk, T. A., Ning, Y., Hsu, M. S., Peng, B., Mains, R. E., Eipper, B. A., et al. (2005). Deletion of peptide amidation enzymatic activity leads to edema and embryonic lethality in the mouse. *Developmental Biology*, 287, 301–313.
- Date, Y., Mondal, M. S., Kageyama, H., Ghamari-Langroudi, M., Cone, R. D., & Nakazato, M. (2010). Neuropeptide W: An anorectic peptide regulated by leptin and metabolic state. *Endocrinology*, 151, 2200–2210.
- Francone, V. P., Ifrum, M. F., Rajagopal, C., Leddy, C. J., Wang, Y., Carson, J. H., et al. (2010). Signaling from the secretory granule to the nucleus: Uhmk1 and PAM. *Molecular Endocrinology*, 24, 1543–1558.

- Grimmelikhuijzen, C. J. P., Hauser, F., Eriksen, F. F., & Williamson, M. (1999). Invertebrate neurohormones and their receptors. *Results and Problems in Cell Differentiation*, *26*, 339–362.
- Gruber, K. A., Fan, W., Akerberg, H., Larhammar, D., Chee, M. J. S., Colmers, W. F., et al. (2009). Neuropeptide Y and gamma-melanocyte stimulating hormone share a common pressor mechanism of action. *Endocrine*, *35*, 312–324.
- Hallberg, M., & Nyberg, F. (2003). Neuropeptide conversion to bioactive fragments—an important pathway in neuromodulation. *Current Protein & Peptide Science*, *4*, 31–44.
- Helke, C. J., Krause, J. E., Mantyh, P. W., Couture, R., & Bannon, M. J. (1990). Diversity in mammalian tachykinin peptidergic neurons: Multiple peptides, receptors, and regulatory mechanisms. *The FASEB Journal*, *4*, 1606–1615.
- Henrich, S., Lindberg, I., Bode, W., & Than, M. E. (2005). Proprotein convertase models based on the crystal structures of furin and kexin: Explanation of their specificity. *Journal of Molecular Biology*, *345*, 211–227.
- Hokfelt, T., Blacker, D., Broberger, C., Herrera-Marschitz, M., & Snyder, G. (2002). Some aspects on the anatomy and function of central cholecystokinin systems. *Pharmacology and Toxicology*, *91*, 382–386.
- Jimenez, E. C., & Olivera, B. M. (2010). Divergent M- and O-superfamily peptides from venom of fish-hunting *Conus pariu*. *Peptides*, *31*, 1678–1683.
- Jimenez-Andrade, J. M., Mantyh, W. G., Bloom, A. P., Xu, H., Vanderah, T. W., & Mantyh, P. W. (2010). A phenotypically restricted set of primary afferent nerve fibers innervate the bone versus skin: Therapeutic opportunity for treating skeletal pain. *Bone*, *46*, 306–313.
- Kalantaridou, S. N., Zoumakis, E., Makrigiannakis, A., Lavasidis, L. G., Vrekoussis, T., &

- Chrousos, G. P. (2010). Corticotropin-releasing hormone, stress and human reproduction: An update. *Journal of Reproductive Immunology*, *85*, 33–39.
- Karhunen, T., Vilim, F. S., Alexeeva, V., Weiss, K. R., & Church, P. J. (2001). Targeting of peptidergic vesicles in cotransmitting terminals. *Journal of Neuroscience*, *21*, RC127.
- Lee, S. N., & Lindberg, I. (2008). 7B2 Prevents unfolding and aggregation of prohormone convertase 2. *Endocrinology*, *149*, 4116–4127.
- Lee, S. N., Peng, B., Desjardins, R., Pintar, J. E., Day, R., & Lindberg, I. (2007). Strain-specific steroidal control of pituitary function. *Journal of Endocrinology*, *192*, 515–525.
- Li, C., Kim, K., & Nelson, L. S. (1999). FMRFamide-related neuropeptide gene family in *Caenorhabditis elegans*. *Brain Research*, *848*, 26–34.
- Mains, R. E., & Eipper, B. A. (1990). The tissue-specific processing of pro-ACTH/endorphin. *Trends in Endocrinology and Metabolism*, *1*, 388–394.
- Mantyh, P. W. (2002). Neurobiology of substance P and the NK1 receptor. *Journal of Clinical Psychiatry*, *63*(Suppl. 11), 6–10.
- McPherson, J., Rivero, G., Baptist, M., Llorente, J., Henderson, G., & Kelly, E. (2010). Mu-opioid receptors: Correlation of agonist efficacy for signalling with ability to activate internalization. *Molecular Pharmacology* (PMID: 20647394).
- Millan, M. J., Dekeyne, A., Gobert, A., Mannoury la Cour, C., Olivier, B., & de Nanteuil, G. (2010). S41744, a dual neurokinin (NK)1 receptor antagonist and serotonin (5-HT) reuptake inhibitor with potential antidepressant properties: A comparison to aprepitant (MK869) and paroxetine. *European Neuropsychopharmacology*, *20*, 599–621.
- Peinado, J. R., Laurent, V., Lee, S. N., Peng, B. W., Pintar, J. E., Steiner, D. F., et al. (2005). Strain-dependent influences on the hypothalamo-pituitary-adrenal axis profoundly affect

- the 7B2 and PC2 null phenotypes. *Endocrinology*, *146*, 3438–3444.
- Prigge, S. T., Mains, R. E., Eipper, B. A., & Amzel, L. M. (2000). New insights into copper monooxygenases and peptide amidation: Structure, mechanism and function. *Cellular and Molecular Life Sciences*, *57*, 1236–1259.
- Rajagopal, C., Stone, K. L., Francone, V. P., Mains, R. E., & Eipper, B. A. (2009). Secretory Granule to the Nucleus: Role of a multiply phosphorylated intrinsically unstructured domain. *Journal of Biological Chemistry*, *284*, 25723–25734.
- Reinscheid, R. K., Nothacker, H. P., Bourson, A., Ardati, A., & Civelli, O. (1995). Orphanin FQ: A neuropeptide that activates an opioidlike G protein-coupled receptor. *Science*, *270*, 792–794.
- Reubi, J. C. (2007). Peptide receptor expression in GEP-NET. *Virchows Archiv*, *451*, S47–S50.
- Sakurai, T., Amemiya, A., Ishii, M., Matsuzaki, I., & Yanagisawa, M. (1998). Orexins and orexin receptors: A family of hypothalamic neuropeptides and G protein-coupled receptors that regulate feeding behavior. *Cell*, *92*, 573–585.
- Sandberg, R., Yasuda, R., Pankratz, D. G., Carter, T. A., Lockhart, D. J., & Barlow, C. (2000). Regional and strain-specific gene expression mapping in the adult mouse brain. *Proceedings of the National Academy of Sciences U.S.A.*, *97*, 11038–11043.
- Sarzi-Puttini, P., Atzeni, F., & Cazzola, M. (2010). Neuroendocrine therapy of fibromyalgia syndrome: An update. *Annals of the New York Academy of Sciences*, *1193*, 91–97.
- Schilling, S., Zeitschel, U., Hoffmann, T., Heiser, U., Cynis, H., Demuth, H. U., et al. (2008). Glutaminyl cyclase inhibition attenuates pyroglutamate A-beta and Alzheimer's disease-like pathology. *Nature Medicine*, *14*, 1106–1111.
- Seidah, N. G., & Chretien, M. (1999). Proprotein and prohormone convertases: A family of

- subtilases generating diverse bioactive polypeptides. *Brain Research*, 848, 45–62.
- Seidah, N. G., Mayer, G., Zaid, A., Rousselet, E., Nassoury, N., Poirier, S., et al. (2008). The activation and physiological functions of the proprotein convertases. *The International Journal of Biochemistry & Cell Biology*, 40, 1111–1125.
- Stephan, A., Wermann, M., von Bohlen, A., Koch, B., Cynis, H., Demuth, H. U., et al. (2009). Mammalian glutaminyl cyclases and their isoenzymes have identical enzymatic characteristics. *The FEBS Journal*, 276, 6522–6536.
- Taghert, P. H., & Veenstra, J. A. (2003). Drosophila neuropeptide signaling. *Advances in Genetics*, 49, 1–65.
- Takanaga, H., Mackenzie, B., & Hediger, M. A. (2004). Sodium-dependent ascorbic acid transporter family SLC23. *Pflugers Archiv*, 447, 677–682.
- Takigami, T., Takeuchi, F., Nakagawa, M., Hase, T., & Tsubaki, M. (2003). Stopped-flow analyses on the reaction of ascorbate with cytochrome b561 purified from bovine chromaffin vesicle membranes. *Biochemistry*, 42, 8110–8118.
- Tekinay, A. B., Nong, Y., Miwa, J. M., Lieberam, I., Ibanez-Tallon, I., Greengard, P., et al. (2009). A role for LYNX2 in anxiety-related behavior. *Proceedings of the National Academy of Sciences U.S.A.*, 106, 4477–4482.
- Terskiy, A., Wannemacher, K. M., Prem, N., Yadav, P. N., Tsai, M., Tian, B., et al. (2007). Search of the human proteome for endomorphin-1 and endomorphin-2 precursor proteins. *Life Sciences*, 81, 1593–1601.
- Wang, S. S., Yan, X. B., Hofman, M. A., Swaab, D. F., & Zhou, J. N. (2010). Increased expression level of corticotropin-releasing hormone in the amygdala and in the hypothalamus in rats exposed to chronic unpredictable mild stress. *Neuroscience Bulletin*,

26, 297–303.

- Westmuckett, A. D., Hoffhines, A. J., Borghei, A., & Moore, K. L. (2008). Early postnatal pulmonary failure and primary hypothyroidism in mice with combined TPST-1 and TPST-2 deficiency. *General and Comparative Endocrinology*, *156*, 145–153.
- Xie, J., Price, M. P., Berger, A. L., & Welsh, M. J. (2002). DRASIC contributes to pH-gated currents in large dorsal root ganglion sensory neurons by forming heteromultimeric channels. *Journal of Neurophysiology*, *87*, 2835–2843.
- Yamaguchi, M., Yamada, Y., Hosokawa, Y., Iwamoto, R., Hoshida, Y., & Matsuzawa, Y. (2010). Long-term suppressive effect of octreotide on progression of metastatic gastrinoma with multiple endocrine neoplasia type 1: Seven-year follow-up. *Internal Medicine*, *49*, 1557–1563.
- Zhou, A., Webb, G., Zhu, X., & Steiner, D. F. (1999). Proteolytic processing in the secretory pathway. *Journal of Biological Chemistry*, *274*, 20745–20748.