

References

- Agulhon, C., Petracicz, J., McMullen, A. B., Sweger, E. J., Minton, S. K., Taves, S. R., et al. (2008). What is the role of astrocyte calcium in neurophysiology? *Neuron*, *59*(6), 932–946.
- Art, J. J., Wu, Y. C., & Fettiplace, R. (1995). The calcium-activated potassium channels of turtle hair cells. *Journal of General Physiology*, *105*(1), 49–72.
- Berridge, M. J., Bootman, M. D., & Roderick, H. L. (2003). Calcium signalling: Dynamics, homeostasis and remodelling. *Nature Reviews Molecular Cell Biology*, *4*(7), 517–529.
- Bobo, R., Bredoux, R., Corvazier, E., Andersen, J. P., Clausen, J. D., Dode, L., et al. (2004). Identification, expression, function, and localization of a novel (sixth) isoform of the human sarco/endoplasmic reticulum Ca²⁺ ATPase 3 gene. *Journal of Biological Chemistry*, *279*(23), 24297–24306.
- Bouchard, R., Pattarini, R., & Geiger, J. D. (2003). Presence and functional significance of presynaptic ryanodine receptors. *Progress in Neurobiology*, *69*(6), 391–418.
- Cornell-Bell, A. H., Finkbeiner, S. M., Cooper, M. S., & Smith, S. J. (1990). Glutamate induces calcium waves in cultured astrocytes: Long-range glial signaling. *Science*, *247*(4941), 470–473.
- Demaurex, N., & Frieden, M. (2003). Measurements of the free luminal ER Ca²⁺ concentration with targeted "chameleon" fluorescent proteins. *Cell Calcium*, *34*(2), 109–119.
- Emptage, N., Bliss, T. V., & Fine, A. (1999). Single synaptic events evoke NMDA receptor-mediated release of calcium from internal stores in hippocampal dendritic spines. *Neuron*, *22*(1), 115–124.
- Feske, S., Gwack, Y., Prakriya, M., Srikanth, S., Puppel, S. H., Tanasa, B., et al. (2006). A

- mutation in Orai1 causes immune deficiency by abrogating CRAC channel function. *Nature*, 441(7090), 179–185.
- Fiacco, T. A., & McCarthy, K. D. (2006). Astrocyte calcium elevations: Properties, propagation, and effects on brain signaling. *Glia*, 54(7), 676–690.
- Filippin, L., Abad, M. C., Gastaldello, S., Magalhaes, P. J., Sandona, D., & Pozzan, T. (2005). Improved strategies for the delivery of GFP-based Ca^{2+} sensors into the mitochondrial matrix. *Cell Calcium*, 37(2), 129–136.
- Giacomello, M., Drago, I., Bortolozzi, M., Scorzeto, M., Gianelle, A., Pizzo, P., et al. (2010). Ca^{2+} hot spots on the mitochondrial surface are generated by Ca^{2+} mobilization from stores, but not by activation of store-operated Ca^{2+} channels. *Molecular Cell*, 38(2), 280–290.
- Golovina, V. A., & Blaustein, M. P. (1997). Spatially and functionally distinct Ca^{2+} stores in sarcoplasmic and endoplasmic reticulum. *Science*, 275(5306), 1643–1648.
- Gordon, G. R. J., Mulligan, S. J., & MacVicar, B. A. (2009). Astrocyte control of blood flow. In V. Parpura & P. G. Haydon (Eds.), *Astrocytes in (patho) physiology of the nervous system*. New York: Springer Science + Business Media.
- Greka, A., Navarro, B., Oancea, E., Duggan, A., & Clapham, D. E. (2003). TRPC5 is a regulator of hippocampal neurite length and growth cone morphology. *Nature Neuroscience*, 6(8), 837–845.
- Halassa, M. M., & Haydon, P. G. (2010). Integrated brain circuits: Astrocytic networks modulate neuronal activity and behavior. *Annual Review of Physiology*, 72, 335–355.
- Huang, G. N., Zeng, W., Kim, J. Y., Yuan, J. P., Han, L., Muallem, S., et al. (2006). STIM1 carboxyl-terminus activates native SOC, I(crac) and TRPC1

- channels. *Nature Cell Biology*, 8(9), 1003–1010.
- Kawabata, S., Tsutsumi, R., Kohara, A., Yamaguchi, T., Nakanishi, S., & Okada, M. (1996). Control of calcium oscillations by phosphorylation of metabotropic glutamate receptors. *Nature*, 383(6595), 89–92.
- Liou, J., Kim, M. L., Heo, W. D., Jones, J. T., Myers, J. W., Ferrell, J. E., Jr., et al. (2005). STIM is a Ca^{2+} sensor essential for Ca^{2+} -store-depletion-triggered Ca^{2+} influx. *Current Biology*, 15(13), 1235–1241.
- Lippincott-Schwartz, J., & Patterson, G. H. (2003). Development and use of fluorescent protein markers in living cells. *Science*, 300(5616), 87–91.
- Llinas, R., Sugimori, M., & Silver, R. B. (1992). Microdomains of high calcium concentration in a presynaptic terminal. *Science*, 256(5057), 677–679.
- Mank, M., & Griesbeck, O. (2008). Genetically encoded calcium indicators. *Chemical Reviews*, 108(5), 1550–1564.
- Marban, E., Rink, T. J., Tsien, R. W., & Tsien, R. Y. (1980). Free calcium in heart muscle at rest and during contraction measured with Ca^{2+} -sensitive microelectrodes. *Nature*, 286(5776), 845–850.
- Miyawaki, A., Llopis, J., Heim, R., McCaffery, J. M., Adams, J. A., Ikura, M., et al. (1997). Fluorescent indicators for Ca^{2+} based on green fluorescent proteins and calmodulin. *Nature*, 388(6645), 882–887.
- Mogami, H., Nakano, K., Tepikin, A. V., & Petersen, O. H. (1997). Ca^{2+} flow via tunnels in polarized cells: Recharging of apical Ca^{2+} stores by focal Ca^{2+} entry through basal membrane patch. *Cell*, 88(1), 49–55.
- Montero, M., Alvarez, J., Scheenen, W. J., Rizzuto, R., Meldolesi, J., & Pozzan, T. (1997). Ca^{2+}

- homeostasis in the endoplasmic reticulum: Coexistence of high and low $[Ca^{2+}]$ subcompartments in intact HeLa cells. *Journal of Cell Biology*, 139(3), 601–611.
- Panatier, A., Theodosis, D. T., Mothet, J. P., Touquet, B., Pollegioni, L., Poulain, D. A., et al. (2006). Glia-derived D-serine controls NMDA receptor activity and synaptic memory. *Cell*, 125(4), 775–784.
- Pascual, O., Casper, K. B., Kubera, C., Zhang, J., Revilla-Sanchez, R., Sul, J. Y., et al. (2005). Astrocytic purinergic signaling coordinates synaptic networks. *Science*, 310(5745), 113–116.
- Permyakov, E. A., & Kretsinger, R. H. (2011). *Calcium binding proteins*. Hoboken, NJ: John Wiley.
- Petzold, G. C., Albeanu, D. F., Sato, T. F., & Murthy, V. N. (2008). Coupling of neural activity to blood flow in olfactory glomeruli is mediated by astrocytic pathways. *Neuron*, 58(6), 897–910.
- Pozzo-Miller, L. D., Pivovarova, N. B., Connor, J. A., Reese, T. S., & Andrews, S. B. (1999). Correlated measurements of free and total intracellular calcium concentration in central nervous system neurons. *Microscopy Research and Technique*, 46(6), 370–379.
- Ridgway, E. B., & Ashley, C. C. (1967). Calcium transients in single muscle fibers. *Biochemical and Biophysical Research Communications*, 29(2), 229–234.
- Ringer, S. (1883). A further contribution regarding the influence of the different constituents of the blood on the contraction of the heart. *Journal of Physiology*, 4(1), 29–42.
- Rizzuto, R., Duchen, M. R., & Pozzan, T. (2004). Flirting in little space: The ER/mitochondria Ca^{2+} liaison. *Science STKE*, 2004(215), re1.
- Rizzuto, R., & Pozzan, T. (2006). Microdomains of intracellular Ca^{2+} : Molecular determinants

- and functional consequences. *Physiological Reviews*, 86(1), 369–408.
- Rogers, M., & Dani, J. A. (1995). Comparison of quantitative calcium flux through NMDA, ATP, and ACh receptor channels. *Biophysical Journal*, 68(2), 501–506.
- Rushton, W. A. H. (1962). *Visual Pigments in Man*. Springfield, IL: Charles C. Thomas.
- Schummers, J., Yu, H., & Sur, M. (2008). Tuned responses of astrocytes and their influence on hemodynamic signals in the visual cortex. *Science*, 320(5883), 1638–1643.
- Shigetomi, E., Kracun, S., Sofroniew, M. V., & Khakh, B. S. (2010). A genetically targeted optical sensor to monitor calcium signals in astrocyte processes. *Nature Neuroscience*, 13(6), 759–766.
- Solovyova, N., & Verkhratsky, A. (2002). Monitoring of free calcium in the neuronal endoplasmic reticulum: An overview of modern approaches. *Journal of Neuroscience Methods*, 122(1), 1–12.
- Strehler, E. E., Filoteo, A. G., Penniston, J. T., & Caride, A. J. (2007). Plasma-membrane Ca^{2+} pumps: Structural diversity as the basis for functional versatility. *Biochemical Society Transactions*, 35(Pt 5), 919–922.
- Takei, K., Stukenbrok, H., Metcalf, A., Mignery, G. A., Sudhof, T. C., Volpe, P., et al. (1992). Ca^{2+} stores in Purkinje neurons: Endoplasmic reticulum subcompartments demonstrated by the heterogeneous distribution of the InsP3 receptor, Ca^{2+} -ATPase, and calsequestrin. *Journal of Neuroscience*, 12(2), 489–505.
- Tsien, R. Y. (1980). New calcium indicators and buffers with high selectivity against magnesium and protons: Design, synthesis, and properties of prototype structures. *Biochemistry*, 19(11), 2396–2404.
- Villa, A., Podini, P., Clegg, D. O., Pozzan, T., & Meldolesi, J. (1991). Intracellular Ca^{2+} stores

- in chicken Purkinje neurons: Differential distribution of the low affinity–high capacity Ca^{2+} binding protein, calsequestrin, of Ca^{2+} ATPase and of the ER luminal protein, Bip. *Journal of Cell Biology*, 113(4), 779–791.
- Wang, X., Lou, N., Xu, Q., Tian, G. F., Peng, W. G., Han, X., et al. (2006). Astrocytic Ca^{2+} signaling evoked by sensory stimulation *in vivo*. *Nature Neuroscience*, 9(6), 816–823.
- Wang, Y., Deng, X., Mancarella, S., Hendron, E., Eguchi, S., Soboloff, J., et al. (2010). The calcium store sensor, STIM1, reciprocally controls Orai and CaV1.2 channels. *Science*, 330(6000), 105–109.
- Yuhas, W. A., & Fuchs, P. A. (1999). Apamin-sensitive, small-conductance, calcium-activated potassium channels mediate cholinergic inhibition of chick auditory hair cells. *Journal of Comparative Physiology A*, 185(5), 455–462.
- Zheng, J. Q. (2000). Turning of nerve growth cones induced by localized increases in intracellular calcium ions. *Nature*, 403(6765), 89–93.
- Zweifach, A., & Lewis, R. S. (1993). Mitogen-regulated Ca^{2+} current of T lymphocytes is activated by depletion of intracellular Ca^{2+} stores. *Proceedings of the National Academy of Sciences of the United States of America*, 90(13), 6295–6299.