

## References

- Acevedo, L., Yu, J., Erdjument-Bromage, H., Miao, R. Q., Kim, J. E., Fulton, D., et al. (2004). A new role for Nogo as a regulator of vascular remodeling. *Nature Medicine*, *17*, 382–388.
- Alexander, M. S., Anderson, K. D., Biering-Sorensen, F., Blight, A. R., Brannon, R., Bryce, T. N., et al. (2009). Outcome measures in spinal cord injury: Recent assessments and recommendations for future directions. *Spinal Cord*, *47*, 582–591.
- Atwal, J. K., Pinkston-Gosse, J., Syken, J., Stawicki, S., Wu, Y., Shatz, C., et al. (2008). PirB is a functional receptor for myelin inhibitors of axonal regeneration. *Science*, *322*, 967–970.
- Bareyre, F. M., Haudenschield, B., & Schwab, M. E. (2002). Long-lasting sprouting and gene expression changes induced by the monoclonal antibody IN-1 in the adult spinal cord. *Journal of Neuroscience*, *22*, 7097–7110.
- Bareyre, F. M., Kerschensteiner, M., Raineteau, O., Mettenleiter, T. C., Weinmann, O., & Schwab, M. E. (2004). The injured spinal cord spontaneously forms a new intraspinal circuit in adult rats. *Nature Neuroscience*, *3*, 269–277.
- Bartsch, U., Bandtlow, C. E., Schnell, L., Bartsch, S., Spillmann, A. A., Rubin, B. P., et al. (1995). Lack of evidence that myelin-associated glycoprotein is a major inhibitor of axonal regeneration in the CNS. *Neuron*, *15*, 1375–1381.
- Benowitz, L. I., & Routtenberg, A. (1997). GAP-43: An intrinsic determinant of neuronal development and plasticity. *Trends in Neuroscience*, *20*, 84–91.
- Bradbury, E. J., Moon, L. D., Popat, R. J., King, V. R., Bennett, G. S., Patel, P. N., et al. (2002). Chondroitinase ABC promotes functional recovery after spinal cord injury. *Nature*, *416*, 636–640.
- Buffo, A., Zagrebelsky, M., Huber, A. B., Skerra, A., Schwab, M. E., Strata, P., et al. (2000).

- Application of neutralizing antibodies against NI-35/250 myelin-associated neurite growth inhibitory proteins to the adult rat cerebellum induces sprouting of uninjured Purkinje cell axons. *Journal of Neuroscience*, 20, 2275–2286.
- Cafferty, W. B., Duffy, P., Huebner, E., & Strittmatter, S. M. (2010). MAG and OMgp synergize with Nogo-A to restrict axonal growth and neurological recovery after spinal cord trauma. *Journal of Neuroscience*, 30, 6825–6837.
- Cajal, R. Y. (1928). *Degeneration and Regeneration of the Nervous System*. New York: Hafner Publishing Co.
- Castro, A. J. (1990). Plasticity in the motor system. In B. Kolb & R. C. Tees (Eds.), *The cerebral cortex of the rat* (pp. 563–588). M.I.T.
- Cheatwood, J. L., Emerick, A. J., Schwab, M. E., & Kartje, G. L. (2008). Nogo-A expression after focal ischemic stroke in the adult rat. *Stroke*, 39, 2091–2098.
- Chen, M. S., Huber, A. B., Van Der Haar, M. E., Frank, M., Schnell, L., Spillmann, A. A., et al. (2000). Nogo-A is a myelin-associated neurite outgrowth inhibitor and an antigen for monoclonal antibody IN-1. *Nature*, 403, 434–439.
- David, S., & Aguayo, A. J. (1981). Axonal elongation into peripheral nervous system ‘bridges’ after central nervous system injury in adult rats. *Science*, 214, 931–933.
- Dimou, L., Schnell, L., Montani, L., Duncan, C., Simonen, M., Schneider, R., et al. (2006). Nogo-A-deficient mice reveal strain-dependent differences in axonal regeneration. *Journal of Neuroscience*, 26, 5591–5603.
- Emerick, A. J., Neafsey, E. J., Schwab, M. E., & Kartje, G. L. (2003). Functional reorganization of the motor cortex in adult rats after cortical lesion and treatment with monoclonal antibody IN-1. *Journal of Neuroscience*, 23, 4826–4830.

- Faulkner, J. R., Herrmann, J. E., Woo, M. J., Tansey, K. E., Doan, M. B., & Sofroniew, M. V. (2004). Reactive astrocytes protect tissue and preserve function after spinal cord injury. *Journal of Neuroscience*, *24*, 2143–2155.
- Fawcett, J. W., Rosser, A. E., & Dunnett, S. B. (2001). Peripheral nerve regeneration: *Brain Damage, Brain Repair*. New York: Oxford University Press.
- Filbin, M. T. (2003). Myelin-associated inhibitors of axonal regeneration in the adult mammalian CNS. *Nature Reviews Neuroscience*, *4*, 703–713.
- Fournier, A. E., Grandpre, T., & Strittmatter, S. M. (2001). Identification of a receptor mediating Nogo-66 inhibition of axonal regeneration. *Nature*, *409*, 341–346.
- Fournier, A. E., Takizawa, B. T., & Strittmatter, S. M. (2003). Rho kinase inhibition enhances axonal regeneration in the injured CNS. *Journal of Neuroscience*, *23*, 1416–1423.
- Gillani, R. L., Tsai, S. Y., Wallace, D. G., O'Brien, T. E., Arhebamen, E., Tole, M., et al. (2010). Cognitive recovery in the aged rat after stroke and anti-Nogo-A immunotherapy. *Behavioral Brain Research*, *208*, 415–424.
- Gonzenbach, R. R., & Schwab, M. E. (2007). Disinhibition of neurite growth to repair the injured adult CNS: Focusing on Nogo. *Cellular and Molecular Life Sciences*, *65*, 161–176.
- GrandPré, T., Li, S., & Strittmatter, S. M. (2002). Nogo-66 receptor antagonist peptide promotes axonal regeneration. *Nature*, *417*, 547–551.
- Grill, R., Murai, K., Blesch, A., Gage, F. H., & Tuszynski, M. H. (1997). Cellular delivery of neurotrophin-3 promotes corticospinal axonal growth and partial functional recovery after spinal cord injury. *Journal of Neuroscience*, *17*, 5560–5572.
- Huber, A. B., Weinmann, O., Brösamle, C., Oertle, T., & Schwab,

- M. E. (2002). Patterns of Nogo mRNA and protein expression in the developing and adult rat and after CNS lesion. *Journal of Neuroscience*, *22*, 3553–3567.
- Joset, A., Dodd, D. A., Haleboua, S., & Schwab, M. E. (2010). Pincher-generated Nogo-A endosomes mediate growth cone collapse and retrograde signaling. *Journal of Cell Biology*, *188*, 271–285.
- Kim, J. E., Li, S., GrandPré, T., Qiu, D., & Strittmatter, S. M. (2003). Axon regeneration in young adult mice lacking Nogo-A/B. *Neuron*, *38*, 187–199.
- Kobayashi, N. R., Fan, D. -P., Giehl, K. M., Bedard, A. M., Wiegand, S. J., & Tetzlaff, W. (1997). BDNF and NT-4/5 prevent atrophy of rat rubrospinal neurons after cervical axotomy, stimulate GAP-43 and T $\alpha$ 1-Tubulin mRNA expression, and promote axonal regeneration. *Journal of Neuroscience*, *17*, 9583–9595.
- Kwon, B. K., Liu, J., Messerer, C., Kobayashi, N. R., McGraw, J., Oschipok, L., et al. (2002). Survival and regeneration of rubrospinal neurons one year after spinal cord injury. *Proceedings of the National Academy of Sciences*, *99*, 3246–3251.
- Lee, J. K., Geoffroy, C. G., Chan, A. F., Tolentino, K. E., Crawford, M. J., Leal, M. A., et al. (2010). Assessing spinal axon regeneration and sprouting in Nogo-, MAG-, and OMgp-deficient mice. *Neuron*, *66*, 663–670.
- Li, S., Liu, B. P., Budel, S., Li, M., Ji, B., Walus, L., et al. (2004). Blockade of Nogo-66, myelin-associated glycoprotein, and oligodendrocyte myelin glycoprotein by soluble Nogo-66 receptor promotes axonal sprouting and recovery after spinal injury. *Journal of Neuroscience*, *24*, 10511–10520.
- Liebscher, T., Schnell, L., Schnell, D., Scholl, J., Schneider, R., Gullo, M., et al. (2005). Nogo-A antibody improves regeneration and locomotion of spinal cord-injured rats. *Annals of*

- Neurology*, 58, 706–719.
- Livesey, F. J., O'Brien, J. A., Li, M., Smith, A. G., Murphy, L. J., & Hunt, S. P. (1997). A Schwann cell mitogen accompanying regeneration of motor neurons. *Nature*, 390, 614–618.
- Markus, T. M., Tsai, S. Y., Bollnow, M. R., Farrer, R. G., O'Brien, T. E., Kindler-Baumann, D. R., et al. (2005). Recovery and brain reorganization after stroke in adult and aged rats. *Annals of Neurology*, 58, 950–953.
- Meiri, K. F., Saffell, J. L., Walsh, F. S., & Doherty, P. (1998). Neurite outgrowth stimulated by neural cell adhesion molecules requires growth-associated protein-43 (GAP-43) function and is associated with GAP-43 phosphorylation in growth cones. *Journal of Neuroscience*, 18, 10429–10437.
- Menei, P., Montero-Menei, C., Whittemore, S. R., Bunge, R. P., & Bunge, M. B. (1998). Schwann cells genetically modified to secrete human BDNF promote enhanced axonal regrowth across transected adult rat spinal cord. *European Journal of Neuroscience*, 10, 607–621.
- Mi, S., Lee, X., Shao, Z., Thill, G., Ji, B., Relton, J., et al. (2004). LINGO-1 is a component of the Nogo-66 receptor/p75 signaling complex. *Nature Neuroscience*, 3, 221–228.
- Montani, L., Gerrits, B., Gehrig, P., Kempf, A., Dimou, L., Wollscheid, B., et al. (2009). Neuronal Nogo-A modulates growth cone motility via Rho-GTP/LIMK1/cofilin in the unlesioned adult nervous system. *Journal of Biological Chemistry*, 284, 10793–10807.
- Oertle, T., & Schwab, M. E. (2003). Nogo and its paRTNers. *Trends in Cell Biology*, 13, 187–194.
- Oertle, T., Van Der Haar, M. E., Bandtlow, C. E., Robeva, A.,

- Burfeind, P., Buss, A., et al. (2003). Nogo-A inhibits neurite outgrowth and cell spreading with three discrete regions. *Journal of Neuroscience*, *23*, 5393–5406.
- Papadopoulos, C. M., Tsai, S. Y., Cheatwood, J. L., Bollnow, M. R., Kolb, B. E., Schwab, M. E., et al. (2006). Dendritic plasticity in the adult rat following middle cerebral artery occlusion and Nogo-a neutralization. *Cerebral Cortex*, *16*, 529–536.
- Papadopoulos, C. M., Tsai, S. -Y., Aslbiei, T., O'Brien, T. E., Schwab, M. E., & Kartje, G. L. (2002). Functional recovery and neuroanatomical plasticity following middle cerebral artery occlusion and IN-1 antibody treatment in the adult rat. *Annals of Neurology*, *51*, 433–441.
- Park, J. B., Yiu, G., Kaneko, S., Wang, J., Chang, J., He, X. L., et al. (2005). A TNF receptor family member, TROY, is a coreceptor with Nogo receptor in mediating the inhibitory activity of myelin inhibitors. *Neuron*, *45*, 345–351.
- Perry, V. H., Brown, M. C., Lunn, E. R., Tree, P., & Gordon, S. (1990). Evidence that very slow wallerian degeneration in C57BL/Ola Mice is an intrinsic property of the peripheral nerve. *European Journal of Neuroscience*, *2*, 802–808.
- Petrinovic, M. M., Duncan, C. S., Bourikas, D., Weinman, O., Montani, L., Schroeter, A., et al. (2010). Neuronal Nogo-A regulates neurite fasciculation, branching and extension in the developing nervous system. *Development*, *137*, 2539–2550.
- Pizzorusso, T., Medini, P., Berardi, N., Chierzi, S., Fawcett, J. W., & Maffei, L. (2002). Reactivation of ocular dominance plasticity in the adult visual cortex. *Science*, *298*, 1248–1251.
- Raineteau, O., & Schwab, M. E. (2001). Plasticity of motor systems after incomplete spinal cord

- injury. *Nature Reviews Neuroscience*, 2, 263–274.
- Schnell, L., Schneider, R., Kolbeck, R., Barde, Y. A., & Schwab, M. E. (1994). Neurotrophin-3 enhances sprouting of corticospinal tract during development and after adult spinal cord lesion. *Nature*, 367, 170–173.
- Schnell, L., & Schwab, M. E. (1990). Axonal regeneration in the rat spinal cord produced by an antibody against myelin-associated neurite growth inhibitors. *Nature*, 343, 269–272.
- Schwab, M. E. (2004). Nogo and axon regeneration. *Current Opinion in Neurobiology*, 14, 118–124.
- Shamash, S., Reichert, F., & Rotshenker, S. (2002). The cytokine network of Wallerian degeneration: Tumor necrosis factor-alpha, interleukin-1-alpha, and interleukin-1-beta. *Journal of Neuroscience*, 22, 3052–3060.
- Shao, Z., Browning, J. L., Leem, X., Scott, M. L., Shulga-Morskaya, S., Allaire, N., et al. (2005). TAJ/TROY, an orphan TNF receptor family member, binds Nogo-66 receptor 1 and regulates axonal regeneration. *Neuron*, 45, 353–359.
- Sicotte, M., Tsatas, O., Jeong, S. Y., Cai, C. -Q., He, Z., & David, S. (2003). Immunization with myelin or recombinant Nogo-66/MAG promotes axon regeneration and sprouting after corticospinal tract lesions in the spinal cord. *Molecular and Cellular Neuroscience*, 23, 251–263.
- Silver, J., & Miller, J. H. (2004). Regeneration beyond the glial scar. *Nature Neuroscience Reviews*, 5, 146–156.
- Simonen, M., Pedersen, V., Weinmann, O., Schnell, L., Buss, A., Ledermann, B., et al. (2003). Systemic deletion of the myelin-associated outgrowth inhibitor Nogo-A improves regenerative and plastic responses after spinal cord injury. *Neuron*, 38, 201–211.

- Tsai, S. Y., Papadopoulos, C. M., Schwab, M. E., & Kartje, G. L. (2011). Delayed anti-Nogo-A therapy improves function after chronic stroke in adult rats. *Stroke*, *42*, 186–190.
- Ughrin, Y. M., Chen, Z. J., & Levine, J. M. (2003). Multiple regions of the NG2 proteoglycan inhibit neurite growth and induce growth cone collapse. *Journal of Neuroscience*, *23*, 175–186.
- Walmsley, A. R., & Mir, A. K. (2007). Targeting the Nogo-A signalling pathway to promote recovery following acute CNS injury. *Current Pharmaceutical Design*, *13*, 2470–2484.
- Wang, K. C., Kim, J. A., Sivasankaran, R., Segal, R., & He, Z. (2002). P75 interacts with the Nogo receptor as a co-receptor for Nogo, MAG and OMgp. *Nature*, *420*, 74–78.
- Wang, K. C., Koprivica, V., Kim, J. A., Sivasankaran, R., Guo, Y., Neve, R. L., et al. (2002). Oligodendrocyte-myelin glycoprotein is a Nogo receptor ligand that inhibits neurite outgrowth. *Nature*, *417*, 941–944.
- Wang, M. S., Fang, G., Culver, D. G., Davis, A. A., Rich, M. M., & Glass, J. D. (2001). The Wld<sup>s</sup> protein protects against axonal degeneration: A model of gene therapy for peripheral neuropathy. *Annals of Neurology*, *50*, 773–779.
- Wiessner, C., Bareyre, F. M., Allegrini, P. R., Mir, A. K., Frenzel, S., Zurini, M., et al. (2003). Anti-Nogo-A antibody infusion 24 hours after experimental stroke improved behavioral outcome and corticospinal plasticity in normotensive and spontaneously hypertensive rats. *Journal of Cerebral Blood Flow and Metabolism*, *23*, 154–165.
- Zheng, B., Ho, C., Li, S., Keirstead, H., Steward, O., & Tessier-Lavigne, M. (2003). Lack of enhanced spinal regeneration in Nogo-deficient mice. *Neuron*, *38*, 213–224.
- Z'Graggen, W. J., Fouad, K., Raineteau, O., Metz, G. A. S., Schwab, M. E., & Kartje, G. L. (2000). Compensatory sprouting and impulse rerouting after unilateral pyramidal tract

lesion in -neonatal rats. *Journal of Neuroscience*, 20, 6561–6569.