

References

- Adler, E., Hoon, M. A., Mueller, K. L., Chandrashekar, J., Ryba, N. J., & Zuker, C. S. (2000). A novel family of mammalian taste receptors. *Cell*, *100*(6), 693–702.
- Bakalyar, H. A., & Reed, R. R. (1990). Identification of a specialized adenylyl cyclase that may mediate odorant detection. *Science*, *250*(4986), 1403–1406.
- Belluscio, L., Gold, G. H., Nemes, A., & Axel, R. (1998). Mice deficient in G(olf) are anosmic. *Neuron*, *20*(1), 69–81.
- Brunet, L. J., Gold, G. H., & Ngai, J. (1996). General anosmia caused by a targeted disruption of the mouse olfactory cyclic nucleotide-gated cation channel. *Neuron*, *17*(4), 681–693.
- Buck, L., & Axel, R. (1991). A novel multigene family may encode odorant receptors: A molecular basis for odor recognition. *Cell*, *65*(1), 175–187.
- Chandrashekar, J., Hoon, M. A., Ryba, N. J., & Zuker, C. S. (2006). The receptors and cells for mammalian taste. *Nature*, *444*(7117), 288–294.
- Chandrashekar, J., Kuhn, C., Oka, Y., Yarmolinsky, D. A., Hummler, E., Ryba, N. J., et al. (2010). The cells and peripheral representation of sodium taste in mice. *Nature*, *464*(7286), 297–301.
- Chandrashekar, J., Yarmolinsky, D., von Buchholtz, L., Oka, Y., Sly, W., Ryba, N. J., et al. (2009). The taste of carbonation. *Science*, *326*(5951), 443–445.
- Dotson, C. D., Vignes, S., Steinle, N. I., & Munger, S. D. (2010). T1R and T2R receptors: The modulation of incretin hormones and potential targets for the treatment of type 2 diabetes mellitus. *Current Opinion in Investigational Drugs*, *11*(4), 447–454.
- Firestein, S. (2001). How the olfactory system makes sense of scents. *Nature*, *413*(6852), 211–218.

- Hoon, M. A., Adler, E., Lindemeier, J., Battey, J. F., Ryba, N. J., & Zuker, C. S. (1999). Putative mammalian taste receptors: A class of taste-specific GPCRs with distinct topographic selectivity. *Cell*, *96*(4), 541–551.
- Jones, D. T., & Reed, R. R. (1989). Golf: An olfactory neuron specific-G protein involved in odorant signal transduction. *Science*, *244*(4906), 790–795.
- Kaupp, U. B., & Seifert, R. (2002). Cyclic nucleotide-gated ion channels. *Physiological Reviews*, *82*(3), 769–824.
- Leybold, B. G., Yu, C. R., Leinders-Zufall, T., Kim, M. M., Zufall, F., & Axel, R. (2002). Altered sexual and social behaviors in *trp2* mutant mice. *Proceedings of the National Academy of Sciences of the United States of America*, *99*(9), 6376–6381.
- Li, X., Staszewski, L., Xu, H., Durick, K., Zoller, M., & Adler, E. (2002). Human receptors for sweet and umami taste. *Proceedings of the National Academy of Sciences of the United States of America*, *99*(7), 4692–4696.
- Liberles, S. D., & Buck, L. B. (2006). A second class of chemosensory receptors in the olfactory epithelium. *Nature*, *442*(7103), 645–650.
- Liman, E. R., Corey, D. P., & Dulac, C. (1999). TRP2: A candidate transduction channel for mammalian pheromone sensory signaling. *Proceedings of the National Academy of Sciences of the United States of America*, *96*(10), 5791–5796.
- Malnic, B., Hirono, J., Sato, T., & Buck, L. B. (1999). Combinatorial receptor codes for odors. *Cell*, *96*(5), 713–723.
- Matsunami, H., Montmayeur, J. P., & Buck, L. B. (2000). A family of candidate taste receptors in human and mouse. *Nature*, *404*(6778), 601–604.
- McLaughlin, S. K., McKinnon, P. J., & Margolskee, R. F. (1992). Gustducin is a taste-cell–

- specific G protein closely related to the transducins. *Nature*, 357(6379), 563–569.
- Mombaerts, P. (2004). Genes and ligands for odorant, vomeronasal and taste receptors. *Nature Reviews Neuroscience*, 5(4), 263–278.
- Munger, S. D., Lane, A. P., Zhong, H., Leinders-Zufall, T., Yau, K. W., Zufall, F., et al. (2001). Central role of the CNGA4 channel subunit in Ca²⁺-calmodulin-dependent odor adaptation. *Science*, 294(5549), 2172–2175.
- Munger, S. D., Leinders-Zufall, T., McDougall, L. M., Cockerham, R. E., Schmid, A., Wandernoth, P., et al. (2010). An olfactory subsystem that detects carbon disulfide and mediates food-related social learning. *Current Biology*, 20(16), 1438–1444.
- Munger, S. D., Leinders-Zufall, T., & Zufall, F. (2009). Subsystem organization of the mammalian sense of smell. *Annual Review of Physiology*, 71, 115–140.
- Nakamura, T., & Gold, G. H. (1987). A cyclic nucleotide-gated conductance in olfactory receptor cilia. *Nature*, 325(6103), 442–444.
- Nelson, G., Chandrashekar, J., Hoon, M. A., Feng, L., Zhao, G., Ryba, N. J., et al. (2002). An amino-acid taste receptor. *Nature*, 416(6877), 199–202.
- Nelson, G., Hoon, M. A., Chandrashekar, J., Zhang, Y., Ryba, N. J., & Zuker, C. S. (2001). Mammalian sweet taste receptors. *Cell*, 106(3), 381–390.
- Perez, C. A., Huang, L., Rong, M., Kozak, J. A., Preuss, A. K., Zhang, H., et al. (2002). A transient receptor potential channel expressed in taste receptor cells. *Nature Neuroscience*, 5(11), 1169–1176.
- Ressler, K. J., Sullivan, S. L., & Buck, L. B. (1993). A zonal organization of odorant receptor gene expression in the olfactory epithelium. *Cell*, 73(3), 597–609.

- Ressler, K. J., Sullivan, S. L., & Buck, L. B. (1994). Information coding in the olfactory system: Evidence for a stereotyped and highly organized epitope map in the olfactory bulb. *Cell*, 79(7), 1245–1255.
- Riviere, S., Challet, L., Fluegge, D., Spehr, M., & Rodriguez, I. (2009). Formyl peptide receptor-like proteins are a novel family of vomeronasal chemosensors. *Nature*, 459(7246), 574–577.
- Serizawa, S., Miyamichi, K., & Sakano, H. (2004). One neuron–one receptor rule in the mouse olfactory system. *Trends in Genetics*, 20(12), 648–653.
- Spehr, M., Gisselmann, G., Poplawski, A., Riffell, J. A., Wetzal, C. H., Zimmer, R. K., et al. (2003). Identification of a testicular odorant receptor mediating human sperm chemotaxis. *Science*, 299(5615), 2054–2058.
- Spehr, M., & Munger, S. D. (2009). Olfactory receptors: G protein-coupled receptors and beyond. *Journal of Neurochemistry*, 109(6), 1570–1583.
- Stephan, A. B., Shum, E. Y., Hirsh, S., Cygnar, K. D., Reisert, J., & Zhao, H. (2009). ANO2 is the ciliary calcium-activated chloride channel that may mediate olfactory amplification. *Proceedings of the National Academy of Sciences of the United States of America*, 106(28), 11776–11781.
- Stowers, L., Holy, T. E., Meister, M., Dulac, C., & Koentges, G. (2002). Loss of sex discrimination and male–male aggression in mice deficient for TRP2. *Science*, 295(5559), 1493–1500.
- Vassar, R., Chao, S. K., Sitcheran, R., Nunez, J. M., Vosshall, L. B., & Axel, R. (1994). Topographic organization of sensory projections to the olfactory bulb. *Cell*, 79(6), 981–991.

- Vassar, R., Ngai, J., & Axel, R. (1993). Spatial segregation of odorant receptor expression in the mammalian olfactory epithelium. *Cell*, *74*(2), 309–318.
- Vigues, S., Dotson, C. D., & Munger, S. D. (2009). The receptor basis of sweet taste in mammals. *Results and Problems in Cell Differentiation*, *47*, 187–202.
- Wong, G. T., Gannon, K. S., & Margolskee, R. F. (1996). Transduction of bitter and sweet taste by gustducin. *Nature*, *381*(6585), 796–800.
- Wong, S. T., Trinh, K., Hacker, B., Chan, G. C., Lowe, G., Gaggar, A., et al. (2000). Disruption of the type III adenylyl cyclase gene leads to peripheral and behavioral anosmia in transgenic mice. *Neuron*, *27*(3), 487–497.
- Young, J. M., Friedman, C., Williams, E. M., Ross, J. A., Tonnes-Priddy, L., & Trask, B. J. (2002). Different evolutionary processes shaped the mouse and human olfactory receptor gene families. *Human Molecular Genetics*, *11*(5), 535–546.
- Zhang, Y., Hoon, M. A., Chandrashekar, J., Mueller, K. L., Cook, B., Wu, D., et al. (2003). Coding of sweet, bitter, and umami tastes: Different receptor cells sharing similar signaling pathways. *Cell*, *112*(3), 293–301.
- Zufall, F., & Leinders-Zufall, T. (2000). The cellular and molecular basis of odor adaptation. *Chemical Senses*, *25*(4), 473–481.
- Zufall, F., & Munger, S. D. (2010). Receptor guanylyl cyclases in mammalian olfactory function. *Molecular and Cellular Biochemistry*, *334*(1–2), 191–197.