

References

- Ahmed, Z. M., Goodyear, R., Riazuddin, S., Lagziel, A., Legan, P. K., Behra, M., et al. (2006). The tip-link antigen, a protein associated with the transduction complex of sensory hair cells, is protocadherin-15. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *26*(26), 7022–7034.
- Avraham, K. B., Hasson, T., Steel, K. P., Kingsley, D. M., & Russell, L. B. (1995). The mouse Snell's waltzer deafness gene encodes an unconventional myosin. *Nature Genetics*, *11*(4), 369–375.
- Beurg, M., Fettiplace, R., Nam, J. H., & Ricci, A. J. (2009). Localization of inner hair cell mechanotransducer channels using high-speed calcium imaging. *Nature Neuroscience*, *12*(5), 553–558.
- Boeda, B., El-Amraoui, A., Bahloul, A., Goodyear, R., Daviet, L., Blanchard, S., et al. (2002). Myosin VIIa, harmonin and cadherin 23, three Usher I gene products that cooperate to shape the sensory hair cell bundle. *EMBO Journal*, *21*(24), 6689–6699.
- Chung, Y. D., Zhu, J., Han, Y., & Kernan, M. J. (2001). *nompA* encodes a PNS-specific ZP domain protein required to connect mechanosensory dendrites to sensory structures. *Neuron*, *29*(2), 415–428.
- Cueva, J. G., Mulholland, A., & Goodman, M. B. (2007). Nanoscale organization of the MEC-4 DEG/ENaC sensory mechanotransduction channel in *Caenorhabditis elegans* touch receptor neurons. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *27*(51), 14089–14098.
- Dumont, R. A., Lins, U., Filoteo, A. G., Penniston, J. T., Kachar, B., & Gillespie, P. G. (2001). Plasma membrane Ca²⁺-ATPase isoform 2a is the PMCA of hair bundles. *Journal of*

- Neuroscience*, 21(14), 5066–5078.
- Ernstrom, G. G., & Chalfie, M. (2002). Genetics of sensory mechanotransduction. *Annual Review of Genetics*, 36, 411–453.
- Fettiplace, R., & Hackney, C. M. (2006). The sensory and motor roles of auditory hair cells. *Nature Reviews. Neuroscience*, 7(1), 19–29.
- Fettiplace, R., & Ricci, A. J. (2003). Adaptation in auditory hair cells. *Current Opinion in Neurobiology*, 13(4), 446–451.
- Gillespie, P. G., & Walker, R. G. (2001). Molecular basis of mechanosensory transduction. *Nature*, 413, 194–202.
- Gillespie, P. G., & Cyr, J. L. (2004). Myosin-1c, the hair cell's adaptation motor. *Annual Review of Physiology*, 66, 521–545.
- Göpfert, M. C., Albert, J. T., Nadrowski, B., & Kamikouchi, A. (2006). Specification of auditory sensitivity by *Drosophila* TRP channels. *Nature Neuroscience*, 9(8), 999–1000.
- Grillet, N., Xiong, W., Reynolds, A., Kazmierczak, P., Sato, T., Lillo, C., et al. (2009). Harmonin mutations cause mechanotransduction defects in cochlear hair cells. *Neuron*, 62(3), 375–387.
- Hackney, C. M., Mahendrasingam, S., Jones, E. M., & Fettiplace, R. (2003). The distribution of calcium buffering proteins in the turtle cochlea. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, 23(11), 4577–4589.
- Heffner, H. E., & Heffner, R. S. (2008). High-frequency hearing. In P. Dallos, D. Oertel & R. Hoy (Eds.), *Handbook of the senses: Audition*. New York, NY: Elsevier. pp. 55–60.
- Heller, S., Bell, A. M., Denis, C. S., Choe, Y., & Hudspeth, A. J. (2002). Parvalbumin 3 is an

- abundant Ca^{2+} buffer in hair cells. *Journal of the Association for Research in Otolaryngology: JARO*, 3(4), 488–498.
- Hill, J. K., Williams, D. E., LeMasurier, M., Dumont, R. A., Strehler, E. E., & Gillespie, P. G. (2006). Splice-site A choice targets plasma-membrane Ca^{2+} -ATPase isoform 2 to hair bundles. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, 26(23), 6172–6180.
- Holt, J. R., Gillespie, S. K., Provance, D. W., Shah, K., Shokat, K. M., Corey, D. P., et al. (2002). A chemical–genetic strategy implicates myosin-1c in adaptation by hair cells. *Cell*, 108(3), 371–381.
- Kachar, B., Parakkal, M., Kurc, M., Zhao, Y., & Gillespie, P. G. (2000). High-resolution structure of hair-cell tip links. *Proceedings of the National Academy of Sciences of the United States of America*, 97(24), 13336–13341.
- Kang, L., Gao, J., Schafer, W. R., Xie, Z., & Xu, X. Z. (2010). C. elegans TRP family protein TRP-4 is a pore-forming subunit of a native mechanotransduction channel. *Neuron*, 67(3), 381–391.
- Kazmierczak, P., Sakaguchi, H., Tokita, J., Wilson-Kubalek, E. M., Milligan, R. A., Muller, U., et al. (2007). Cadherin 23 and protocadherin 15 interact to form tip-link filaments in sensory hair cells. *Nature*, 449(7158), 87–91.
- Kernan, M., Cowan, D., & Zuker, C. (1994). Genetic dissection of mechanosensory transduction: Mechanoreception-defective mutations of *Drosophila*. *Neuron*, 12(6), 1195–1206.
- Kloda, A., & Martinac, B. (2002). Common evolutionary origins of mechanosensitive ion channels in Archaea, Bacteria and cell-walled Eukarya. *Archaea (Vancouver, B.C.)*, 1(1), 35–44.

- Kros, C. J., Marcotti, W., van Netten, S. M., Self, T. J., Libby, R. T., Brown, S. D., et al. (2002). Reduced climbing and increased slipping adaptation in cochlear hair cells of mice with *Myo7a* mutations. *Nature Neuroscience*, *5*, 41–47.
- Kung, C. (2005). A possible unifying principle for mechanosensation. *Nature*, *436*(7051), 647–654.
- Lumpkin, E. A., & Hudspeth, A. J. (1998). Regulation of free Ca^{2+} concentration in hair-cell stereocilia. *Journal of Neuroscience*, *18*(16), 6300–6318.
- O'Hagan, R., Chalfie, M., & Goodman, M. B. (2005). The MEC-4 DEG/ENaC channel of *Caenorhabditis elegans* touch receptor neurons transduces mechanical signals. *Nature Neuroscience*, *8*(1), 43–50.
- Rzadzinska, A. K., Schneider, M. E., Davies, C., Riordan, G. P., & Kachar, B. (2004). An actin molecular treadmill and myosins maintain stereocilia functional architecture and self-renewal. *Journal of Cell Biology*, *164*(6), 887–897.
- Sakaguchi, H., Tokita, J., Naoz, M., Bowen-Pope, D., Gov, N. S., & Kachar, B. (2008). Dynamic compartmentalization of protein tyrosine phosphatase receptor Q at the proximal end of stereocilia: Implication of myosin VI-based transport. *Cell Motility and the Cytoskeleton*, *65*(7), 528–538.
- Salles, F. T., Merritt, R. C. J., Manor, U., Dougherty, G. W., Sousa, A. D., Moore, J. E., et al. (2009). Myosin IIIa boosts elongation of stereocilia by transporting espin 1 to the plus ends of actin filaments. *Nature Cell Biology*, *11*(4), 443–450.
- Senften, M., Schwander, M., Kazmierczak, P., Lillo, C., Shin, J. B., Hasson, T., et al. (2006). Physical and functional interaction between protocadherin 15 and myosin VIIa in

- mechanosensory hair cells. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, 26(7), 2060–2071.
- Shin, J. B., Longo-Guess, C. M., Gagnon, L. H., Saylor, K. W., Dumont, R. A., Spinelli, K. J., et al. (2010). The R109H variant of fascin-2, a developmentally regulated actin crosslinker in hair-cell stereocilia, underlies early-onset hearing loss of DBA/2J mice. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, 30(29), 9683–9694.
- Sidi, S., Friedrich, R. W., & Nicolson, T. (2003). NompC TRP channel required for vertebrate sensory hair cell mechanotransduction. *Science*, 301(5629), 96–99.
- Siemens, J., Kazmierczak, P., Reynolds, A., Sticker, M., Littlewood-Evans, A., & Muller, U. (2002). The Usher syndrome proteins cadherin 23 and harmonin form a complex by means of PDZ-domain interactions. *Proceedings of the National Academy of Sciences of the United States of America*, 99(23), 14946–14951.
- Siemens, J., Lillo, C., Dumont, R. A., Reynolds, A., Williams, D. S., Gillespie, P. G., et al. (2004). Cadherin 23 is a component of the tip link in hair-cell stereocilia. *Nature*, 428, 950–955.
- Sukharev, S., & Corey, D. P. (2004). Mechanosensitive channels: Multiplicity of families and gating paradigms. *Science's STKE [Electronic Resource]: Signal Transduction Knowledge Environment*, 2004(219), re4.
- Tobin, D., Madsen, D., Kahn-Kirby, A., Peckol, E., Moulder, G., Barstead, R., et al. (2002). Combinatorial expression of TRPV channel proteins defines their sensory functions and subcellular localization in *C. elegans* neurons. *Neuron*, 35(2), 307–318.
- Todi, S. V., Franke, J. D., Kiehart, D. P., & Eberl, D. F. (2005). Myosin VIIA defects, which underlie the Usher 1B syndrome in humans, lead to deafness in *Drosophila*. *Current*

Biology: CB, 15(9), 862–868.

- Verpy, E., Leibovici, M., Zwaenepoel, I., Liu, X. Z., Gal, A., Salem, N., et al. (2000). A defect in harmonin, a PDZ domain-containing protein expressed in the inner ear sensory hair cells, underlies usher syndrome type 1C. *Nature Genetics*, 26(1), 51–55.
- Walker, R. G., Willingham, A. T., & Zuker, C. S. (2000). A *Drosophila* mechanosensory transduction channel. *Science*, 287(5461), 2229–2234.
- Walsh, T., Walsh, V., Vreugde, S., Hertzano, R., Shahin, H., Haika, S., et al. (2002). From flies' eyes to our ears: Mutations in a human class III myosin cause progressive nonsyndromic hearing loss DFNB30. *Proceedings of the National Academy of Sciences of the United States of America*, 99(11), 7518–7523.
- Wang, V. Y., Hassan, B. A., Bellen, H. J., & Zoghbi, H. Y. (2002). *Drosophila* atonal fully rescues the phenotype of Math1 null mice: New functions evolve in new cellular contexts. *Current Biology: CB*, 12(18), 1611–1616.
- Weil, D., El-Amraoui, A., Masmoudi, S., Mustapha, M., Kikkawa, Y., Laine, S., et al. (2003). Usher syndrome type I G (USH1G) is caused by mutations in the gene encoding SANS, a protein that associates with the USH1C protein, harmonin. *Human Molecular Genetics*, 12(5), 463–471.
- Yan, J., Pan, L., Chen, X., Wu, L., & Zhang, M. (2010). The structure of the harmonin/sans complex reveals an unexpected interaction mode of the two Usher syndrome proteins. *Proceedings of the National Academy of Sciences of the United States of America*, 107(9), 4040–4045.
- Zheng, L., Sekerkova, G., Vranich, K., Tilney, L. G., Mugnaini, E., & Bartles, J. R. (2000). The deaf jerker mouse has a mutation in the gene encoding the espin actin-bundling proteins

of

hair

cell

stereocilia and lacks espins. *Cell*, 102(3), 377–385.