

Proteomics



SUMMARY

Proteomics is the study of protein products from genes using a global approach. There are many techniques in biotechnology for studying proteins. Just as DNA can be separated on agarose based on size using DNA's negative charge, polyacrylamide gel electrophoresis (PAGE) is used to separate proteins based on size. However, not all proteins have an overall negative charge. Sodium dodecyl sulfate (SDS) is added to protein samples, and with high heat the protein denatures and is coated with a negative charge. The proteins are visualized within the gel using stains. Some proteins are overall positively charged, whereas others are neutral or negatively charged. Isoelectric focusing uses a pH gradient in the electrophoresis setup to separate proteins based on native charge alone. Two-dimensional PAGE (2D-PAGE) increases the resolving power of gel electrophoresis for whole cell protein extracts by separating based on native charge first and then on size.

Western blotting is a useful tool to identify proteins. Protein samples are subjected to SDS-PAGE first and then transferred to a membrane, such as nitrocellulose. The membrane is treated with antibodies that recognize specific epitopes on the protein of interest. A secondary antibody that recognizes the primary antibody is then added to the membrane. The secondary has a conjugated detection system so that the location of binding can be determined, often by using chemiluminescence or chromogenic substances.

Proteins can be separated, purified, identified, and quantified using a process called high-pressure liquid chromatography (HPLC). Proteins are dissolved in a solvent and then forced through a column packed with a solid material. When the sample exits the column on the opposite end, it is collected for further analysis. The solid material present in the column dictates the property of proteins used in the separation. Size exclusion chromatography uses a solid material containing porous beads to separate based on size. Reverse phase HPLC separates proteins based on the hydrophilic or hydrophobic nature. The solid material in ion-exchange HPLC contains charged molecules, which are then used to separate proteins based on charge. Affinity chromatography uses a solid material containing a molecule, such as an antibody, that binds to a specific protein. Changing conditions in the liquid phase passing over the solid phase can disrupt the association between protein and antibody, for example, and cause the protein to elute out of the column almost entirely together. The molecules exiting the column are detected by monitoring light refraction, UV light absorption, or fluorescence, or are detected radiochemically or electrochemically.

Proteases are used to degrade proteins. Some proteases are specific to certain sequences, much like restriction enzymes specific to DNA sequences. Other proteases are less specific and are capable of completely digesting proteins to amino acid monomers. Proteases are particularly important for use in mass spectrometry (MS), which is used to measure the mass of ions in a sample. Since proteins are very large molecules, they must first be prepared by breaking them into smaller pieces for MS. In matrix-assisted laser desorption-ionization (MALDI), proteins are embedded onto a solid matrix and then ionized to determine molecular mass. In electrospray ionization (ESI), proteins are dissolved in a liquid that is then evaporated under high heat, which breaks apart the protein into ions. The mass of these ions is then determined by MS. The more futuristic surface-enhanced laser desorption-ionization (SELDI) relies on proteins attached to a solid metal bar, and they are then ionized for analysis by MS. SELDI could be used one day to detect changes in proteins within a droplet of body fluid. MS is sensitive enough to detect changes in phosphorylation and glycosylation and has broad implications in medical testing and diagnosis. The multidimensional protein identification technique (MuDPIT) involves fragmenting a mixture of proteins, which are then identified by MS. Prior to MS, in MuDPIT, a 2D LC microcapillary column separated the protein fragments first by charge and then by size and hydrophobicity. Samples up to 3000 different proteins can be evaluated on one column and one MS experiment.



Proteins can also be quantified using MS. SILAC, which is an acronym for Stable Isotope Labeling by Amino acids in Cell culture, uses cells that are grown with or without stable isotope-labeled amino acids. Heavier isotopes increase the mass of all proteins. When the two samples are compared by MS, two peaks are observed. The ratio of the heavy isotope with the normal protein determines the change in protein level.

Additionally, variations on currently available MS techniques are used to determine the sequence of proteins in shorter pieces of polypeptides. The concentration of proteins can also be quantified by inclusion of amino acids labeled with heavy isotopes. A comparison of MS data yields relative changes in protein concentration between specific experimental conditions.

Tagging proteins with various labels is helpful in isolation and identification of specific proteins. Generally, the tag is fused with the gene that encodes the protein of interest so that they are translated together. Various tags are currently being used in research. The most common is the polyhistidine tag (His_{6X}), which is made up of six tandem histidine residues. This tag could be attached to either the C-terminus or the N-terminus end of the protein. The His_{6X} tag readily binds divalent metal ions and, therefore, is easily isolated and separated from a mixture of proteins through affinity chromatography. Other commonly used short tags include FLAG, which binds to anti-FLAG antibodies for separation, and Strep, which binds to avidin or streptavidin present on beads in a column. Proteins can also be tagged with larger molecules, such as protein A, glutathione-S-transferase (GST), and maltose-binding protein (MBP). The separating mechanism is usually achieved through antibodies to the tag, binding to glutathione or maltose, respectively. Addition of a protease cleavage site allows the tag to be removed once the protein of interest is purified.

Phage display libraries are bacteriophage expressing foreign proteins on their outer surfaces. The gene for the protein of interest is fused with a gene from the bacteriophage, which then expresses the protein once it infects a host. These phage libraries are helpful to identify proteins that bind to cell surface receptors or to specific RNA molecules.

The two-hybrid yeast system allows the study of interactions between proteins. For example, hormone interaction with receptors is studied using this system. In this system, the interaction of the two proteins elicits a response from a reporter gene that is used to monitor the interaction. GAL4 is a transcriptional activator that contains two domains in order to function: a DNA-binding domain (DBD) and an activation domain (AD). One protein of interest is fused to the DBD, and the second protein of interest is fused to the AD. All fusion occurs at the genetic level, so it is important to ensure that they are fused in frame. For transcription of a reporter gene to be activated, the DBD and AD must be associated. If the fused proteins of interest interact, they draw together the DBD and AD to form a complete GAL4, which then activates transcription of the reporter gene.

The two-hybrid yeast system examines protein interactions in the nucleus, but it is also possible to examine protein interactions in the cytoplasm using a technique called co-immunoprecipitation. In this technique, an antibody to a protein (or if an antibody is unavailable, a protein tag) is used to sequester the protein, along with anything bound to it. Analysis occurs via isolation and separation using SDS-PAGE followed by MS to determine the sequence.

Protein expression can be monitored by comparing protein extracts from two different conditions and then subjecting them to protein arrays. Antigen capture immunoassay uses antibodies bound to a solid surface. Protein samples are labeled with fluorescent dyes and then mixed with the array. If the protein binds to the array, it fluoresces. Multiple conditions are compared using a different dye for each condition. In direct immunoassay, or reverse-phase array, the proteins are bound to a solid surface and then probed with a specific fluorescently-labeled antibody. In this experiment, multiple samples, perhaps from different patients, can be spotted onto the same solid support and probed simultaneously for the presence of a specific protein. Other protein arrays exist that eliminate the need for antibodies,



which can often cross-react, giving unwanted results. More recently, protein arrays have been used to bind small hairpin RNA molecules.

With more and more refined techniques available for the study of biological systems, it is becoming increasingly possible to study all of metabolism for a cell on a global scale. Metabolomics describes the small molecules and intermediates of metabolism that exist within a system at one particular time. Since metabolism is dynamic, only snapshots of metabolism can be obtained through the use of nuclear magnetic resonance (NMR), MS, and HPLC. Since metabolites can affect the pigments, scents, flavors, and nutrient content of plants, studying the plant metabolome is especially prudent. These studies could enhance flavors and yield fresher produce.



Case Study Proteomic Mapping of the Human Mitochondrial Intermembrane Space in Live Cells via Ratiometric APEX Tagging

Victoria Hung et al. (2014). *Molecular Cell* 55, 332–341.

Complete lists of the proteins present within specific regions of cells are difficult to obtain and often rely on traditional mass spectrometry–based proteomics that require high yields and purity of components that can actually be isolated. The authors of this paper developed a mapping method for specific proteomes in living cells. APEX is an engineered ascorbate peroxidase that is targeted to the subcellular compartment. APEX then tags all proteins in the surrounding area. The tagged proteins are then isolated and identified by MS.

APEX has been used to map the proteome of the human mitochondrial matrix. The authors of this article used APEX to map the proteome of the mitochondrial intermembrane space (IMS). Labeling the IMS proteins has the potential to generate background results because it is possible for APEX to label some cytoplasmic proteins near the mitochondria. The authors developed a SILAC-based strategy to ensure APEX labeling was specific to the IMS.

How did the authors ensure that APEX was correctly localized to the IMS?

The authors genetically engineered APEX to the IMS by fusing it to the peptide leader sequence of an IMS native protein. Confirmation of the localization was through electron microscopy.

Some of the biotinylated proteins on the imaging results showed a diffuse pattern that extended beyond the mitochondria. What might account for this phenomenon? How did the authors determine which possibility was correct?

During the labeling procedures, there was a one-minute window in which the biotin-phenoxyl radicals produced in the labeling process diffused beyond the mitochondria. Another possibility is that the labeled products were able to diffuse away from the mitochondria in that same one-minute window. The authors developed Western blot assays to determine that, likely, relocation of biotin-labeled proteins and other molecules occurred rather than the diffusion of radicals.

The initial proteomic experiments yielded high cytosolic background signals. What method did the authors develop to map the IMS proteome with more specificity? What was the outcome?

The authors developed a ratiometric approach that measured the ratio of biotinylation extent of two constructs: one construct inside the IMS (IMS-APEX) and the other on the outside of the region of interest (cytosolic APEX). This ratio-based process works because one would expect an IMS protein to be more extensively labeled by the IMS-APEX than by the cytosolic APEX. Three SILAC cultures were used: heavy culture for IMS-APEX, medium culture for cytosolic APEX, and the negative control light culture containing no APEX. This method detected two major populations of biotinylated proteins. One population was enriched IMS proteins. The second population contained mostly nonmitochondrial proteins. These results were consistent with expectations.

Did the authors find any new mitochondrial proteins within the IMS? If so, how did they ensure these signals were not false positives? How did they characterize these proteins?

The results showed that 87% of the proteins identified in the experiment were previously identified as mitochondrial proteins. Sixteen proteins had no previous mitochondrial annotation and could either be false positive or newly discovered IMS proteins. The authors examined six of these proteins through fluorescence imaging and found overlap within the mitochondria. Three other proteins were investigated through Western blot analysis. Three other proteins were found not to be associated with the mitochondria. After considering the proteins with no previous mitochondria annotation, the authors raised the specificity of their experiment to 94%.

In terms of specificity, is it possible that IMS-APEX could label membrane-bound proteins that protrude into the IMS?

By using complexes with known associations to various mitochondrial subcompartments, the authors determined that only those proteins that are exposed within the IMS are detected as part of the IMS proteome using their method. Additionally, the authors examined the actual sites of labeling on the proteins. They sequenced 47 of their 127 identified IMS proteins. Of these 47, 14 were transmembrane IMM proteins. This number indicates that membrane proteins are also labeled but only if they are exposed within the subcompartment of the organelle.

Were there any proteins of particular interest in the IMS proteome that warranted further analysis? If so, which one(s)?

Yes. The authors identified and validated a total of nine newly discovered mitochondrial proteins. These proteins included three that are associated with the endoplasmic reticulum, a lipid transporter, a protein involved in regulation of mitosis, an autophagy receptor, and proteins with unknown function. Of particular interest were the proteins that associate with the endoplasmic reticulum; a protein called MICU1, which is involved in calcium binding and transport; and a protein that is dual-localized to the mitochondrial matrix and IMS.

The authors developed an interesting technique to map the proteome of mitochondrial subcompartments. Their first attempt yielded a high number of false positive and diffuse results. In their refined method, the authors were able to increase the specificity and identified multiple previously annotated proteins as well as some proteins with no previously known mitochondrial annotation. Limitations of this technique include low coverage, which is likely due to stereochemistry and accessibility. The authors also cite the inability to determine the exact amino acid on the protein that is labeled by APEX and imperfect specificity as limitations. Lastly, dual-localized proteins are potentially missed with this methodology. Despite the limitations, their method will be a powerful tool for determining the proteomes within specific cellular and organelle subcompartments. Their method is performed in living cells and certainly has the advantage of less manipulation of the organelles and cells, which could produce artifacts.

Proteomic Mapping of the Human Mitochondrial Intermembrane Space in Live Cells via Ratiometric APEX Tagging

Victoria Hung,¹ Peng Zou,^{1,5} Hyun-Woo Rhee,^{1,2} Namrata D. Udeshi,³ Valentin Cracan,⁴ Tanya Svinikina,³ Steven A. Carr,³ Vamsi K. Mootha,^{3,4} and Alice Y. Ting^{1,3,*}

¹Department of Chemistry, Massachusetts Institute of Technology, Cambridge, MA 02139, USA

²Ulsan National Institute of Science and Technology (UNIST), Ulsan 689-798, Korea

³Broad Institute of MIT and Harvard, Cambridge, MA 02142, USA

⁴Department of Molecular Biology, Howard Hughes Medical Institute, Massachusetts General Hospital, Harvard Medical School, Boston, MA 02114, USA

⁵Present address: Department of Chemistry and Chemical Biology, Harvard University, Cambridge, MA 02138, USA

*Correspondence: ating@mit.edu

<http://dx.doi.org/10.1016/j.molcel.2014.06.003>

SUMMARY

Obtaining complete protein inventories for subcellular regions is a challenge that often limits our understanding of cellular function, especially for regions that are impossible to purify and are therefore inaccessible to traditional proteomic analysis. We recently developed a method to map proteomes in living cells with an engineered peroxidase (APEX) that bypasses the need for organellar purification when applied to membrane-bound compartments; however, it was insufficiently specific when applied to unbounded regions that allow APEX-generated radicals to escape. Here, we combine APEX technology with a SILAC-based ratiometric tagging strategy to substantially reduce unwanted background and achieve nanometer spatial resolution. This is applied to map the proteome of the mitochondrial intermembrane space (IMS), which can freely exchange small molecules with the cytosol. Our IMS proteome of 127 proteins has >94% specificity and includes nine newly discovered mitochondrial proteins. This approach will enable scientists to map proteomes of cellular regions that were previously inaccessible.

INTRODUCTION

Cell biologists strive to obtain complete protein lists for the subcellular regions they are studying, but using traditional mass spectrometry (MS)-based proteomics, this is possible only for cellular compartments that can be isolated in high yield and purity. Because many cellular regions are difficult or impossible to purify, their proteomes are unknown or incompletely known. To address this challenge, we developed a method to map specific proteomes in living cells using an engineered ascorbate peroxidase (APEX) (Martell et al., 2012; Rhee et al., 2013). As shown in Figure 1A, APEX is first genetically targeted to the

cellular compartment of interest. Then, upon addition of the small molecule biotin-phenol and hydrogen peroxide (H₂O₂), APEX covalently tags surrounding endogenous proteins with the biotin-phenoxyl radical oxidation product during a 1 min window. Subsequently, cells are lysed, and biotinylated proteins are isolated with streptavidin beads and identified by MS.

Previously, we used APEX to map the proteome of the human mitochondrial matrix (Rhee et al., 2013). In the present study, we focused on the proteome of the mitochondrial intermembrane space (IMS), which lies between the inner mitochondrial membrane (IMM) and outer mitochondrial membrane (OMM) (Figure 1A). The IMS cannot be purified by traditional approaches such as density centrifugation. This compartment is associated with many essential functions, including apoptosis, protein import and folding, and reactive oxygen species detoxification (Herrmann and Riemer, 2010), yet it is much less well-understood than the mitochondrial matrix.

In addition to its biological importance, the IMS presents a major technical challenge for the APEX methodology. Unlike the mitochondrial matrix, the IMS is not fully membrane-enclosed. Although the IMM blocks the passage of APEX-generated biotin-phenoxyl radicals into the matrix (Rhee et al., 2013), the OMM contains porins that allow free exchange of molecules <5 kDa between the cytosol and IMS (Colombini, 1979). Thus, IMS-targeted APEX could potentially biotinylate cytosolic proteins outside the mitochondria, giving unwanted background (Figure 1A). Here, we develop a stable isotope labeling by amino acids in cell culture (SILAC)-based ratiometric tagging strategy that effectively excludes cytosolic proteins and produces a highly specific proteomic map of the human IMS.

RESULTS

IMS-APEX Labeling in Cells Characterized by Imaging and Western Blotting

We targeted APEX to the IMS of HEK 293T cells by fusing it to the 68 amino acid leader sequence of the native IMS serine β -lactamase-like protein LACTB (Polianskyte et al., 2009) and confirmed correct localization by electron microscopy (EM)




Figure 1. APEX-Based Proteomic Mapping Scheme and Characterization of IMS-APEX Labeling

(A) Scheme. APEX (green Pac-Man) is targeted to the intermembrane space (IMS) of HEK 293T cells by genetic fusion to the leader sequence of the IMS protein LACTB (Polianskyte et al., 2009). The IMS and matrix subcompartments of the mitochondrion are labeled in blue, in addition to the outer mitochondrial membrane (OMM) and inner mitochondrial membrane (IMM). To initiate proteomic tagging, H₂O₂ is added to live cells for 1 min in the presence of biotin-phenol (red B = biotin). APEX catalyzes the generation of biotin-phenoxyl radicals, which tag proximal endogenous proteins. Cells are then lysed, and tagged proteins are purified with streptavidin beads and identified by mass spectrometry. Due to the porosity of the OMM (porins allow passage of molecules <5 kDa [Colombini, 1979]), IMS-APEX can tag some cytosolic proteins outside mitochondria, giving unwanted background. This study reports a ratiometric method to eliminate this background. The enlarged box at right shows one possible structure of the biotin-phenol adduct with a tyrosine side chain (Tyr). Other adduct structures, including with other amino acids, are likely to be formed as well.

(B) Electron microscopic (EM) characterization of IMS-APEX localization. HEK 293T cells expressing IMS-APEX were fixed and then overlaid with a solution of diaminobenzidine (DAB) and H₂O₂. APEX catalyzes the oxidation of DAB to generate a locally deposited DAB polymer (Martell et al., 2012). After staining the polymer with electron-dense OsO₄, the sample was imaged by EM. For comparison, APEX targeted to the mitochondrial matrix (Rhee et al., 2013) is shown on the right. Scale bars, 100 nm.

(C) Streptavidin blot analysis of endogenous proteins tagged by IMS-APEX. Samples were transfected with IMS-APEX or cytosolic APEX-NES (NES = nuclear export sequence) and labeled for 1 min as in (A). Afterwards, cells were lysed, run on a gel, and analyzed by streptavidin blotting (left) and Ponceau staining (right). Negative controls are shown with biotin-phenol (BP) omitted (lanes 2 and 5) or H₂O₂ omitted (lanes 3 and 6). Lane 7 shows untransfected cells. The band near 80 kDa contains the endogenously biotinylated proteins 3-methylcrotonyl-CoA carboxylase and propionyl-CoA carboxylase (Chapman-Smith and Cronan, 1999). (D) Fluorescence imaging of IMS-APEX labeling in cells. HEK 293T cells were transfected with IMS-APEX, matrix-APEX, or APEX-NES and labeled live as in (A). Cells were then fixed and stained with neutravidin to visualize biotinylated proteins and anti-V5 or anti-Flag antibody to visualize APEX localization. The anti-V5/Flag channel is not normalized. DIC, differential interference contrast image. AF647 and AF488 are Alexa Fluor dyes. Scale bars, 20 μm.

(Martell et al., 2012) (Figure 1B). Next, transfected cells were labeled live with H₂O₂ for 1 min in the presence of preincubated biotin-phenol, then lysed and analyzed by gel electrophoresis and streptavidin blot to detect biotinylated proteins. Figure 1C shows that IMS-APEX biotinylates many proteins in a banding pattern that differs from that produced by a cytosolic APEX variant. This suggests that IMS-APEX and cytosolic APEX tag different endogenous proteomes, as expected.

Biotin labeling by IMS-APEX was also assessed by imaging. After live-cell labeling for 1 min, samples were fixed and stained with neutravidin-Alexa-Fluor-647 to visualize biotinylated proteins. Figure 1D shows a diffuse biotin staining pattern that extends well beyond mitochondria, even though IMS-APEX is cleanly localized to mitochondria (Figures 1B and S1A). In contrast, the localization of mitochondrial matrix-APEX overlaps tightly with the proteins it biotinylates (Figure 1D, second row).

This suggests that, in the case of IMS-APEX labeling, either the APEX-generated biotin-phenoxyl radicals diffuse far beyond the OMM (through porins), and/or that molecules biotinylated by IMS-APEX subsequently diffuse away from mitochondria during the 1 min labeling window to give a diffuse pattern.

To distinguish between these two possibilities, we developed a western blot assay. We harvested proteins biotinylated by IMS-APEX using streptavidin beads and probed this enriched material for the presence of both IMS and cytosolic markers. [Figures S1B–S1E](#) show that this protocol enriches the endogenous IMS protein apoptosis-inducing factor (AIF) in addition to a cotransfected IMS marker protein (LACTB-myc-YFP). A cotransfected cytosolic marker (mCherry-Flag-LpIA) is enriched to a much lesser extent, indicating much lower biotinylation by IMS-APEX. In the converse experiment, we found that cytosol-localized APEX strongly biotinylates the Flag-tagged cytosolic marker but much more weakly biotinylates the two IMS proteins. These results suggest that the APEX labeling radius is indeed limited, and the diffuse neutravidin staining observed in [Figure 1D](#) may result from relocalization of biotin-tagged proteins and other molecules, rather than long-range diffusion of the biotin-phenoxyl radical. To more rigorously investigate the spatial specificity of labeling, however, we proceeded to a MS proteomic experiment.

Initial Proteomic Experiment Produces High Cytosolic Background

For our first attempt, we used an experimental setup identical to that of our APEX mitochondrial matrix study ([Rhee et al., 2013](#)). The experimental samples were grown in SILAC media ([Ong et al., 2002](#)) containing heavy isotopes of arginine and lysine, whereas the control samples were grown in light SILAC media ([Figure S2A](#)). More than 4,000 proteins were detected by MS in each of two independent replicates. Protein SILAC ratios are plotted in histograms in [Figures S2D and S2E](#). Encouragingly, true positive and false positive analyses show that the right ends of the histograms (proteins with high heavy/light, or H/L, SILAC ratios) are enriched for known IMS proteins, whereas the left ends (low H/L) contain predominantly nonmitochondrial proteins.

We produced a list of 362 entries by retaining only proteins with high H/L in both replicates (row 1 of [Figure 2D](#)). The mitochondrial specificity was calculated as the fraction of proteins in this list with prior mitochondrial annotation (40%). Depth of coverage was determined by generating a list of 75 well-established IMS proteins (IMS gold+ list, tab 1 of [Table S2](#)) and calculating the fraction of this list present in our proteome (69%). Both specificity and coverage values are much lower than those previously obtained in our mitochondrial matrix study (>94% and ~85%, respectively) ([Rhee et al., 2013](#)). We found that by raising the H/L cut-off (row 2 of [Figure 2D](#)), we could increase specificity to 58% at the expense of coverage (49%). These numbers indicate that this data set cannot be used to generate an IMS proteome of acceptable quality.

Examining the data more closely, we found that high H/L proteins include cytosolic proteins that reside near mitochondria, such as kinesin and microtubule-associated proteins. We interpret this to mean that (1) the porosity of the OMM, as feared, permits IMS-APEX-generated biotin-phenoxyl radicals to escape

the IMS and biotinylate cytosolic proteins, and (2) SILAC ratio magnitude reflects not only proximity to APEX but likely other factors as well, such as steric accessibility and surface tyrosine count, explaining why some detected cytosolic proteins have higher H/L ratios than bona fide IMS proteins. Given these factors, it was necessary to develop a different approach to using APEX for proteomic mapping with high spatial specificity.

Ratiometric APEX Tagging Yields a High-Quality IMS Proteome

In the field of fluorescent reporters, absolute fluorescence intensity in a cell can be affected by many factors other than the specific parameter of interest (e.g., phosphorylation or Ca^{2+} concentration), such as local variations in reporter concentration ([Mehta and Zhang, 2011](#)). To address this problem, many reporters are designed to be ratiometric, such that the emission ratio of two fluorophores reads out the desired parameter and is independent of other factors such as reporter concentration ([Grynkiewicz et al., 1985](#); [Ting et al., 2001](#)). We sought to apply a similar concept to APEX tagging ([Figure 2A](#)). Instead of only measuring the extent of a protein's biotinylation by IMS-APEX (via its H/L SILAC ratio), we sought to measure the ratio of its biotinylation extent by two different APEX constructs: one inside the region of interest (e.g., IMS-APEX) and one outside the region of interest (e.g., cytosolic APEX). This ratio-based strategy would address both of the problems described above. First, even if biotin-phenoxyl radicals escape beyond the OMM, it should be possible to distinguish IMS-resident proteins from cytosolic proteins outside mitochondria by the ratio of their biotinylation extents by IMS-APEX versus cytosolic APEX. This is because an IMS-resident protein should be more extensively labeled by IMS-APEX than by cytosolic APEX, and the reverse should be true for a cytosol-resident protein. Second, this ratio should reflect *only* proximity to the IMS (or cytosol) and no other factor such as steric accessibility because such factors should equally affect IMS-APEX labeling and cytosolic APEX labeling of the same protein and therefore be cancelled out.

To implement this ratiometric approach, we established three rather than two SILAC cultures: a heavy culture (H) that was labeled with IMS-APEX, a medium culture (M) that was labeled with a cytosolic APEX construct (APEX-NES, where NES is a nuclear export signal), and a light culture (L) that was a negative control with APEX omitted ([Figure 2B](#)). While H/L and M/L ratios reflect biotinylation extent by IMS-APEX and APEX-NES, respectively, the H/M ratio reflects the extent to which a protein is preferentially biotinylated by IMS-APEX versus APEX-NES.

More than 4,800 proteins were detected by MS, each associated with three SILAC ratios: H/L, M/L, and H/M. In [Figure 2C](#), each detected protein is plotted by its $\log_2(\text{H/L})$ ratio (reflecting extent of biotinylation by IMS-APEX) on the y axis and by its $\log_2(\text{M/L})$ ratio (reflecting extent of biotinylation by APEX-NES) on the x axis. Two major populations are visible: one with a negative slope in the top left quadrant and one with a slightly positive slope in the top right quadrant. The first population is enriched in known IMS proteins (from our IMS gold+ list, tab 1 of [Table S2](#)), which are colored in green, and the second population contains mostly proteins that lack mitochondrial annotation, which are colored in red. This result is consistent with




Figure 2. Ratiometric APEX Tagging Strategy Improves Spatial Specificity and Produces a High-Quality IMS Proteome

(A) Because the OMM is permeable to molecules <5 kDa, including the biotin-phenoxyl radical, some cytosolic proteins will be tagged by IMS-APEX. However, it is possible to distinguish IMS proteins (e.g., protein 1, represented by the black box) from cytosolic proteins (e.g., protein 2) by comparing each protein's extent of biotinylation by IMS-APEX (left) versus cytosolic APEX (APEX-NES) (right). For example, protein 1 should be more strongly biotinylated by IMS-APEX than by APEX-NES, regardless of its steric accessibility or surface tyrosine count. Conversely, protein 2 should be tagged more extensively by APEX-NES than by IMS-APEX. Red coloring represents endogenous proteins biotinylated by APEX.

(B) Three-state SILAC experimental setup. Three HEK 293T cultures were treated identically with biotin-phenol and H₂O₂ for 1 min, but the heavy culture expressed IMS-APEX, the medium culture expressed APEX-NES, and the light culture was untransfected. After labeling, the three lysate samples were combined and processed together as shown. For each protein, the H/L SILAC ratio reflects the extent of its biotinylation by IMS-APEX. The M/L SILAC ratio reflects the extent of its biotinylation by APEX-NES. The H/M SILAC ratio reflects the ratio of that protein's biotinylation by IMS-APEX versus APEX-NES.

(C) Scatter plot showing H/L ratio plotted against M/L ratio for 99.96% of the 4,868 proteins identified by MS (inset shows all proteins, including the few with very low SILAC ratios). Proteins previously known to be IMS-exposed are colored green in the plot (i.e., true positives, defined as members of our IMS gold+ list [tab 1 of Table S2]). Proteins without previous mitochondrial annotation are colored red (false positives). SILAC cut-offs used in row 4 of (D) are shown by the dashed lines. See "Scatter plot analysis" in the Supplemental Experimental Procedures and tab 5 of Table S1 for details.

(D) Table showing IMS proteome size, specificity, and coverage derived from different data sets and different SILAC cut-offs (truncated to three significant figures). Specificity refers to mitochondrial specificity, i.e., the percentage of the proteome that has mitochondrial annotation in GOCC (Ashburner et al., 2000), Mitocarta (Pagliarini et al., 2008), our previous mitochondrial matrix proteome (Rhee et al., 2013), or the literature. Coverage refers to the percentage of our IMS gold+ list (tab 1 of Table S2) that is detected in the proteome. Condition 5 (bottom row, colored blue) was used to obtain the final IMS proteome (shown in tab 1 of Table S1). Rep1 and Rep2 are the two different replicates of the two-state SILAC experiment described in Figure S2A.

our experimental design because the top left quadrant contains proteins with higher H/L ratios and lower M/L ratios, i.e., high H/M ratios, reflecting greater biotinylation by IMS-APEX than by APEX-NES.

To filter these data and define our IMS proteome, we calculated the optimal SILAC ratio cut-offs that maximize the difference between the true positive rate and the false positive rate (Figure S2F). Interestingly, the application of a cut-off based on




Figure 3. Characterization of IMS Proteome Specificity

(A) Bar graph showing the enrichment of mitochondrial proteins as well as IMS-exposed proteins in the IMS proteome. The first two columns show the percentage of proteins, in the entire human proteome and in the IMS proteome, respectively, with prior mitochondrial annotation. The second two columns show the percentage of proteins with potential IMS exposure (IMS, IMM, or OMM annotation). *mito proteins refers to the 579 mitochondrial proteins with annotated submitochondrial localization. See tab 4 of Table S2 for details.

(B) Subunits of the TOM/TIM/PAM mitochondrial protein import complex (Bolender et al., 2008; Gebert et al., 2011; Neupert and Herrmann, 2007) detected in the IMS proteome (left) and in our previous mitochondrial matrix proteome (right) (Rhee et al., 2013). See tab 5 of Table S2 for details.

(C) Imaging analysis of six mitochondrial orphans identified in this study. After transient transfection, proteins were detected by anti-V5 staining in COS-7 cells and compared to a mitochondrial GFP marker. Regions of overlap are colored yellow in the “merge” row. Quantitation of overlap from ≥ 10 cells for each protein is given beneath each image set. A positive control construct (IMS-APEX) gave $89.3\% \pm 6.3\%$ mitochondrial overlap, while negative control constructs (P4HB-V5 and APEX-NES) gave $40.3\% \pm 9.9\%$ and $27.0\% \pm 9.0\%$ mitochondrial overlap, respectively (data not shown). Scale bars, 10 μm . Note that imaging experiments for individual orphans were performed separately, rather than in parallel, but are shown together here. The error for each percentage is defined as one sample standard deviation with Bessel’s correction.

(D) Western blot detection of three mitochondrial orphans identified in this study, in purified mouse liver mitochondria. WTL is whole tissue lysate. Protein molecular weights are 110 kDa (NBR1), 65 kDa (FKBP10), and 50 kDa (CDC25C). Control blots are shown for a mitochondrial matrix protein (ATP5B, 51 kDa), which becomes enriched as mitochondrial purity increases, and an ER protein (calreticulin, 48 kDa), which becomes de-enriched.

H/M alone produced a list of 237 proteins whose mitochondrial specificity was 72% and whose coverage of known IMS proteins was 68% (row 3 of Figure 2D). These values are already much improved compared to those obtained using our original two-state SILAC approach. When we filtered the data using H/L ratio as well to exclude proteins without biotinylation signal above background, the mitochondrial specificity increased to 80%, and the coverage dropped by only 1% to 67%. For our final IMS proteomic list of 127 proteins (row 5 of Figure 2D and tab 1 of Table S1), we retained proteins with high H/M from the three-state experiment and high H/L from both replicates of our two-state SILAC experiments. The mitochondrial specificity of this list was 87%, and its coverage of known IMS proteins was

65%. These numbers are dramatically improved compared to those obtained with IMS-APEX labeling alone and approach the values for our previous matrix proteome (Rhee et al., 2013).

Characterizing the Specificity of the IMS Proteome

Figure 3A shows that 87% of the proteins in the IMS proteome have prior mitochondrial annotation. Sixteen proteins, or 13%, do not have previous connections to mitochondria (mitochondrial “orphans” in tab 2 of Table S2). These could be false positives or newly discovered mitochondrial proteins. We performed fluorescence imaging of six of these proteins and observed partial or complete overlap with a mitochondrial GFP marker (Figure 3C). Three additional proteins were detected by western

blot analysis of purified mouse mitochondria (Figure 3D). These data raise the actual mitochondrial specificity of the IMS proteome from 87% to 94%. Notably, we also performed imaging and/or western blot analysis of three additional mitochondrial orphans (RCN2, P4HB, and CCSMST1) but did not obtain evidence of mitochondrial localization. These proteins could represent erroneous detections, or they could be present in such low amounts in mitochondria that our assays do not have sufficient sensitivity to detect them.

To analyze the specificity of the proteomic list for IMS proteins in particular, as opposed to proteins in other regions of mitochondria, we first considered the fact that IMS-APEX could potentially tag any IMS-exposed protein, including OMM and IMM proteins. We therefore calculated the fraction of proteins in the IMS proteome with IMS, OMM, or IMM annotation as opposed to matrix annotation. This was 99%, compared to 74% for all mitochondrial proteins that have submitochondrial annotation (Figure 3A). Our IMS proteome is therefore enriched in potentially IMS-exposed proteins.

We also analyzed IMS specificity by examining the components of the TOM/TIM/PAM mitochondrial protein import complex, whose subunits span multiple mitochondrial subcompartments (Bolender et al., 2008; Gebert et al., 2011; Neupert and Herrmann, 2007). Figure 3B shows that only IMS-exposed subunits of this complex are detected in the IMS proteome. This is strikingly complementary to our detection of matrix-exposed subunits in our previous matrix proteome (Rhee et al., 2013). We performed a similar analysis of Complexes I–V of the electron transport chain (Figure S3A). Most of the subunits we detected in the IMS proteome are known to be IMS-exposed. One exception was F6 of Complex V (ATP synthase), which was highly enriched in the IMS proteome and absent from our previous matrix proteome (Rhee et al., 2013). Follow-up EM analysis (Figure S3B) showed F6-APEX2 staining in the matrix only, suggesting that our detection of endogenous F6 in the IMS is erroneous or a small population of F6 exists in the IMS that is not visible by EM.

Finally, IMS specificity can be examined via the biotinylated peptides that we detected, which reveal the exact site of labeling by the biotin-phenoxyl radical. Such peptides were identified and sequenced for 47 of our 127 IMS proteins (tab 3 of Table S1). Of these 47 proteins, 14 are transmembrane IMM proteins with known or predicted topologies. The 24 biotinylated peptides derived from these 14 proteins all map to IMS-exposed rather than matrix-exposed regions (column S in tab 3 of Table S1). Three of these 14 transmembrane IMM proteins (APC1, NDUFB6, and UQCRCQ) had biotinylated peptides detected in the matrix proteomic experiment as well (Rhee et al., 2013) (column T in tab 3 of Table S1). The three sites biotinylated by matrix-APEX map exclusively to matrix-exposed sides (Figure S3C). Our observations reinforce the notion that biotin-phenoxyl radicals do not cross the IMM and suggest that APEX labeling can be used to help unravel the topology of membrane proteins within living cells.

Analysis of Specific Proteins within the IMS Proteome

The IMS proteome provides insights and raises numerous hypotheses, particularly when combined with our previous mito-

chondrial matrix proteome. First, as described above, we have identified nine newly discovered mitochondrial proteins that we validated by imaging or western blotting. These include three endoplasmic reticulum (ER)-associated proteins (FKBP10, MGST3, and STX17), a lipid transport protein (APOO), a mitotic regulator (CDC25C), an autophagy receptor (NBR1), and three proteins of unknown function (A8MTT3, C1orf163, and CCDC127). We were especially intrigued by the ER-annotated proteins because of the great interest in mitochondria-ER communication and contact sites (Kornmann and Walter, 2010). One of our mitochondrial orphans, syntaxin 17 (Stx17), is a transmembrane SNARE-type protein that has been observed at mitochondria-ER contact sites during autophagy (Hamasaki et al., 2013). However, its localization under basal conditions is unclear. Hamasaki et al. observed Stx17 solely on ER membranes (Hamasaki et al., 2013), but Itakura et al. reported cytosolic and mitochondrial pools as well (Itakura et al., 2012). To further examine the localization of Stx17 and reconcile our data with previous literature, we performed EM imaging of an APEX2 (S.S. Lam, J.D. Martell, and A.Y.T., unpublished data) fusion to Stx17. DAB stain was clearly observed on the OMM facing the cytosol and at mitochondria-ER contact sites (Figure 4A). This result supports partial mitochondrial localization of Stx17, with the loop between transmembrane domains contacting the IMS, as shown in Figure 4B.

Second, the IMS proteome gives insight into mitochondrial proteins whose submitochondrial localizations are unknown or under debate. For example, our data indicate that 21 proteins without prior submitochondrial localization data have IMS-exposed regions (tab 3 of Table S2). Furthermore, our data shed light on MICU1, a calcium-binding protein that regulates the function of the mitochondrial calcium uniporter (MCU), the channel in the IMM that allows mitochondria to uptake calcium from the cytosol (Perocchi et al., 2010). The mechanism by which MICU1 regulates MCU is unclear partly because of conflicting evidence regarding MICU1's submitochondrial localization. One study used protease accessibility assays on purified mitochondria to show that MICU1 resides in the IMS (Csordás et al., 2013), where it is able to sense changes in cytosolic Ca^{2+} levels. However, another study used live cell fluorescence imaging to conclude that MICU1 resides in the matrix (Hoffman et al., 2013), which is shielded from cytosolic Ca^{2+} changes by the IMM. Mallilankaraman et al. have proposed a mechanistic model for MICU1 regulation that relies on its residence in the matrix (Mallilankaraman et al., 2012), and this model conflicts with that of Csordás et al., which assumes that MICU1 resides in the IMS (Csordás et al., 2013). These differing conclusions reflect the imperfect nature of the assays used to determine MICU1 submitochondrial localization. For example, these assays can be affected by lysis of purified mitochondria or overexpression and tag interference with MICU1 function.

In contrast, our proteomic data report on endogenous, untagged proteins in intact mitochondria within living cells. MICU1 was detected in our IMS proteome, but not in our matrix proteome (Rhee et al., 2013) (Figure 4C). Interestingly, MICU2, a related protein with nonredundant function (Kamer and Mootha, 2014), was also detected in the IMS proteome but not in the matrix proteome (Rhee et al., 2013). Therefore, the IMS and matrix proteomic data together are consistent with




Figure 4. Localization Analysis of Stx17 and MICU1

(A) EM imaging of syntaxin 17 (Stx17) with an APEX2 tag (S.S. Lam, J.D. Martell, and A.Y.T., unpublished data) fused to its N terminus. HEK 293T were transduced with APEX2-Stx17 lentivirus, then processed as in Figure 1B. DAB staining (arrowheads) is observed at junctions between mitochondria (M) and endoplasmic reticulum (ER) tubules. Scale bars, 200 nm.

(B) Cartoon illustrating possible arrangement of Stx17 molecules at starred mitochondria-ER contact site in (A). Stx17 is a SNARE-type protein with two transmembrane domains and N and C termini that face the cytosol (Itakura et al., 2012). Based on the DAB staining pattern in (A), we propose that Stx17 resides in both OMM and ER membranes with its N terminus facing the cytosol. For the OMM pool, the hairpin loop joining the two transmembrane domains would contact the IMS, explaining the detection of Stx17 in our IMS proteome.

(C) Known components of the mitochondrial calcium uniporter (MCU) complex are depicted at left (Sancak et al., 2013). In the table at right, for each protein of this complex, the corresponding IMS proteome (three-state experiment) and matrix proteome (replicate 1) SILAC values are listed. n.d. indicates not detected. Gray-shaded values were below the cut-offs for inclusion in IMS or matrix proteomes. Red-shaded values were above the cut-offs for inclusion in the IMS proteome. See tab 7 of Table S2 for details.

MICU1 and MICU2 localization in the IMS. These observations will help to develop a mechanistic model for its regulatory behavior.

Third, a set of hypotheses emerge when we cross our IMS proteome with our previous matrix proteome (Rhee et al., 2013). As discussed above, APEX-generated biotin-phenoxyl radicals do not cross the IMM. Hence, a protein that is detected in both the IMS and matrix proteomes is likely to be either a dual-localized protein or a transmembrane IMM protein. Thirty-three proteins were found in both proteomic lists (tab 6 of Table S2), and of these, 19 are known to be transmembrane IMM proteins (e.g., SCO1 [Horg et al., 2005] and LETM1 [Tamai et al., 2008]). One protein, HSPE1, is known to be dual-localized (Samali et al., 1999). The remaining 13 proteins are potentially undescribed transmembrane IMM or IMS/matrix dual-localized proteins that warrant further investigation.

DISCUSSION

APEX is a powerful technology for tagging endogenous proteomes in living cells, but a major question raised by our previous study, which mapped the mitochondrial matrix proteome (Rhee

et al., 2013), is whether or not this method has the spatial specificity to map cellular regions that are not membrane-enclosed. We have addressed this question by mapping the proteome of the mitochondrial IMS. Unlike the matrix, the IMS is “leaky” because the outer mitochondrial membrane contains porins that allow free passage of molecules <5 kDa in size (Colombini, 1979). APEX-generated biotin-phenoxyl radicals (0.362 kDa) could thus potentially diffuse out of the IMS to give unwanted tagging of cytosolic proteins.

In this study, we showed that the original approach cannot produce a sufficiently specific map of the IMS because APEX-generated radicals tag cytosolic proteins just outside mitochondria and produce significant background signal. Therefore, we developed a different methodology based on ratiometric tagging of endogenous proteins by APEX in the IMS versus APEX in the cytosol, followed by data filtering based on ratio magnitude. This approach was able to produce a high-quality IMS proteome of 127 proteins with >94% specificity and 65% coverage. We note that our ratiometric approach has some resemblance to previous approaches (Dunkley et al., 2004; Foster et al., 2006) used to improve specificity of proteomic mapping of purified ER and Golgi.

Though demonstrated only for the IMS here, we expect the ratiometric APEX tagging approach to be generalizable to other cellular regions as well. This technique should always be used with a quantitative proteomic method such as SILAC or iTRAQ (isobaric tag for relative and absolute quantitation). Our study shows that although biotin-phenoxy radical concentrations fall off rapidly, proteins that are tens to hundreds of nanometers away (e.g., beyond the OMM, which is ~5 nm thick) may still be detectably biotinylated; hence, in the absence of an impermeable membrane barrier, the mere presence of a biotin label is not evidence of close proximity to APEX. The ratio of two experimental samples, one with APEX in the region or macromolecular complex of interest, and the other with APEX outside the region or complex (as we did here, with IMS-APEX and APEX-NES), best reflects proximity to APEX.

The methodology has some important limitations. One is that coverage is not high. We suspect that the main factor limiting coverage is steric accessibility of modifiable amino acid side chains (e.g., tyrosine) to the biotin-phenoxy radical. Since labeling is performed in live cells, macromolecular complexes are still intact, which will fundamentally limit the accessibility of sterically hindered proteins to the biotin probe. We are also interested in exploring new labeling chemistries that can target a broader array of side chains.

Second, because the APEX radical chemistry results in chemically heterogeneous adducts of unknown masses, we currently have limited ability to determine the exact residues tagged by the biotin-phenol probe. This information could have great value, for instance for determining the topology of membrane proteins. Improvements in the labeling chemistry and development of peptide-enrichment strategies should allow us to map biotinylation sites much more extensively than is currently possible.

A third limitation to the method is that specificity, though high in this case, is not perfect. For our IMS proteome, seven proteins were identified that we and others could not validate as true mitochondrial proteins. They may be present in the IMS in very low amounts. Alternatively, since five of these are known to be ER or secretory pathway proteins (RCN2, P4HB, NUCB2, TXNDC12, and LAMC1), and the IMS and ER may be physically linked (Chandra et al., 1998), perhaps biotin-phenoxy radicals generated in the IMS leak into the ER to tag endogenous proteins there. Our ratiometric SILAC experiment with APEX-NES would not have excluded this background, but a future experiment with IMS-APEX labeling ratioed against APEX-ER labeling should be able to.

Finally, our approach has the problem of potentially missing dual-localized proteins. For example, a protein that has both an IMS and a cytosolic population would be filtered out based on low H/M ratio. We did generate a separate table of 33 proteins that have high H/L ratios, but only moderate H/M ratios (tab 2 of Table S1); however, these would need to be screened one by one to determine if both IMS and cytosolic pools truly exist (or if they are transmembrane OMM proteins).

Recently, Vögtle et al. mapped the proteome of the yeast IMS by a different approach (Vögtle et al., 2012). Purified mitochondria were treated with the apoptotic factor Bax to trigger permeabilization of the outer mitochondrial membrane. SILAC was then used to determine which IMS proteins were lost following

this treatment. Compared to our approach, the main disadvantages are the reliance on purified mitochondria (which can introduce artifacts, such as the background efflux of matrix proteins observed over time in this study), the lack of detection of membrane-associated proteins that do not dissociate after Bax treatment (for example, an analogous study on mammalian mitochondria would fail to detect MICU1 because it remains tightly associated with MCU in the IMM [Sancak et al., 2013]), and the inability to generalize the approach to species that do not utilize Bax-type signaling. Correspondingly, their IMS proteome of 49 proteins is considerably smaller than our IMS proteome of 127 proteins. Of their 49 yeast proteins, 35 have human orthologs. Of these, 18 were detected in our IMS proteome (column Y in tab 1 of Table S1). For the 17 proteins we did not detect, two are not expressed in HEK 293T (Sultan et al., 2008), and three are likely to be expressed at very low levels (Beck et al., 2011; Sultan et al., 2008). The others may be undetected because they are sterically shielded from biotinylation in the context of living cells, or they are not localized to the IMS in human cells.

The human IMS proteome produced by this study should have value for mitochondrial biologists. We identified 16 proteins not previously associated with mitochondria and confirmed by follow-up imaging and western blotting of purified mitochondria that nine are indeed mitochondrial (Figures 3C and 3D). One of these, Stx17, was observed by EM imaging to be at mitochondria-ER contact sites (Figure 4A). In addition, we assigned 21 mitochondrial proteins not previously known to be IMS-localized to the IMS (tab 3 of Table S2). Finally, our data suggest that a key regulator of mitochondrial calcium uptake, MICU1, resides in the IMS (Figure 4C). Our detection of endogenous MICU1 in the IMS helps to explain how it responds to cytosolic calcium levels to control the activity of the mitochondrial calcium uniporter MCU.

From a methodological perspective, our study establishes that APEX can be used to map cellular proteomes, even in the absence of a tight membrane barrier, with a spatial resolution on the order of ~5 nm (the distance between proteins inside the IMS and proteins just outside the OMM). With the ratiometric tagging approach, it will be interesting and valuable in future studies to map cellular proteomes that are even less understood and more challenging than the IMS, such as the outer mitochondrial membrane and mitochondria-ER contact sites.

EXPERIMENTAL PROCEDURES

SILAC Labeling and Biotinylation of the IMS Proteome

Figure S2A and Figure 2B show the experimental configurations for the two-state and three-state SILAC experiments, respectively. On day 0, early passage HEK 293T cells were seeded into three T25 flasks and cultured in SILAC media to metabolically label their proteomes with heavy, medium, or light isotopes of lysine and arginine (Ong et al., 2002). The media consisted of different isotopes of L-arginine and L-lysine in 10% dialyzed fetal bovine serum (Sigma), penicillin, streptomycin, glutamine, and 4.5 g/l glucose in DMEM (Caisson Laboratories). The first flask was cultured in light SILAC media, which contained L-arginine (Arg0) and L-lysine (Lys0) (Sigma). The second flask was cultured in heavy SILAC media, which contained L-arginine-¹³C₆, ¹⁵N₄ (Arg10) and L-lysine-¹³C₆, ¹⁵N₂ (Lys8) (Sigma). The third flask was cultured in medium SILAC media, which contained L-arginine [¹³C₆]HCl (Arg6) and L-lysine-4,4,5,5-*d*₄ (Lys4) (Sigma). The cells were split into fresh SILAC media every 2 days before they became fully confluent. On day 4, the cultures were expanded into T75

flasks. After two more passages, the cultures were expanded into T150 flasks on day 10 as follows: three T150 flasks in heavy SILAC media (labeled H1, H2, and H3), one T150 flask in medium SILAC media (labeled M3), and four T150 flasks in light SILAC media (labeled L1, L2, L3, and L4).

On day 11, the H1, H2, H3, L2, L3, and L4 cultures were transfected with 15 μ g IMS-APEX plasmid using 120 μ l Lipofectamine 2000 (Invitrogen) in 30 ml heavy or light DMEM without serum or antibiotics. The M3 flask was similarly transfected with 15 μ g APEX-NES using medium DMEM. After 4 hr, the transfection solution was replaced with fresh SILAC media, and the cells were allowed to recover for 24 hr.

Cells were labeled on day 12. The H1, H2, M3, L1, L3, and L4 cultures were incubated with 500 μ M biotin-phenol in 40 ml SILAC media for 30 min at 37°C. Then, H₂O₂ was added to a final concentration of 1 mM for 1 min at room temperature, after which the probe/H₂O₂ solution was replaced with 15 ml of “quencher solution” (10 mM sodium ascorbate, 10 mM sodium azide, and 5 mM Trolox in Dulbecco’s Phosphate Buffered Saline [DPBS]). All samples, including the L2 flask (negative control with biotin-phenol and H₂O₂ omitted), were washed twice with the quencher solution, twice with DPBS, and once more with the quencher solution. Each wash volume was 15 ml. The cells were then collected in 5 ml of quencher solution by gentle pipetting and pelleted at 500 \times g for 3 min at room temperature. After discarding the supernatant, the cell pellet was stored at –80°C overnight.

Cell pellets were thawed on ice and then lysed with 800 μ l of freshly-prepared, ice-cold RIPA lysis buffer (50 mM Tris-HCl [pH 7.4], 150 mM NaCl, 0.1% SDS, 0.5% sodium deoxycholate, 1% Triton X-100, 1 \times protease inhibitor cocktail (Sigma Aldrich catalog no. P8849), 1 mM PMSF (phenylmethylsulfonyl fluoride), 10 mM sodium azide, 10 mM sodium ascorbate, and 5 mM Trolox) by gentle pipetting. The lysates were centrifuged at 13,000 rpm for 10 min at 4°C, and the protein concentrations of the clarified lysates were measured using a Pierce 660 nm Protein Assay kit, with freshly made bovine serum albumin solutions as standards.

Streptavidin Enrichment and Elution of Biotinylated Proteins

The table under “In-gel digestion of biotinylated proteins, extraction, liquid chromatography, and mass spectrometry” in the [Supplemental Experimental Procedures](#) summarizes how the lysates for each SILAC culture were combined. The two-state Replicate 1 sample consisted of H1 and L1 mixed in a 1:1 ratio (by protein concentration). The two-state Replicate 2 sample consisted of H2 and L2 mixed in a 1:1 ratio. The three-state sample consisted of H3, M3, and L3 mixed in a 1:1:1 ratio. Each sample had a total of at least 2.6 mg protein (at 2–4 mg/ml protein concentration). In addition, L4 lysate was spiked in at 5% (by protein concentration) to each sample to facilitate quantitation of isotope ratios.

Streptavidin-coated magnetic beads (Pierce) were washed twice with RIPA lysis buffer. Each sample was mixed with 500 μ l of streptavidin bead slurry. The suspensions were gently rotated at room temperature for 1 hr to bind biotinylated proteins. The flowthrough after enrichment was removed and stored at 4°C, and the beads were washed with 2 \times 1 ml RIPA lysis buffer, 1 ml of 2 M urea in 10 mM Tris-HCl (pH 8.0), and again with 2 \times 1 ml RIPA lysis buffer. These denaturing washes are important to disrupt protein-protein interactions and ensure enrichment only of proteins directly biotinylated by APEX. Biotinylated proteins were then eluted by heating the beads at 95°C for 5 min in 60 μ l 1 \times NuPAGE LDS Sample Buffer (Invitrogen) supplemented with 20 mM DTT and 2 mM biotin. A second round of enrichment was completed on the flowthrough from the three-state SILAC experiment (following the same protocol) because the first round of enrichment did not yield sufficient protein.

Further processing of the biotinylated proteins for mass spectrometry and all other experimental procedures are described in the [Supplemental Experimental Procedures](#).

SUPPLEMENTAL INFORMATION

Supplemental Information includes Supplemental Experimental Procedures, three figures, and two tables and can be found with this article online at <http://dx.doi.org/10.1016/j.molcel.2014.06.003>.

AUTHOR CONTRIBUTIONS

P.Z. and H.-W.R. contributed equally to this work. V.H., P.Z., H.-W.R., N.D.U., S.A.C., V.K.M., and A.Y.T. designed the experiments. V.H., P.Z., and H.-W.R. characterized IMS-APEX labeling and prepared the proteomic samples. N.D.U., T.S., and S.A.C. processed the proteomic samples and performed mass spectrometry. V.H., P.Z., and H.-W.R. performed initial analysis on the proteomic data. V.C. and V.K.M. generated purified mitochondria samples. V.H. did all other experiments and final analysis. V.H. and A.Y.T. wrote the paper. All authors edited the paper.

ACKNOWLEDGEMENTS

We thank J. Martell, E. Vasile (Koch Institute Microscopy Core Facility), and N. Watson (Whitehead Institute Keck Microscopy Facility) for help with EM sample preparation. E. Vasile and N. Watson also acquired the EM images. We thank N. Mizushima for assistance with interpretation of EM images, T. Hashimoto and S. Calvo for advice on data analysis, and H.B. Fraser and J. Martell for assistance with manuscript editing. J. Litke helped clone some of the mitochondrial orphan constructs. V.H. was supported by a National Science Foundation Graduate Research Fellowship. Funding was provided by the NIH (DP1 OD003961 to A.Y.T.), Dreyfus Foundation (A.Y.T.), American Chemical Society (A.Y.T.), and the Howard Hughes Medical Institute Collaborative Initiative Award (A.Y.T. and S.A.C.). V.K.M. is an Investigator of the Howard Hughes Medical Institute. Massachusetts Institute of Technology has submitted a patent application related to this work.

Received: February 27, 2014

Revised: April 21, 2014

Accepted: May 20, 2014

Published: July 3, 2014

REFERENCES

- Ashburner, M., Ball, C.A., Blake, J.A., Botstein, D., Butler, H., Cherry, J.M., Davis, A.P., Dolinski, K., Dwight, S.S., Eppig, J.T., et al.; The Gene Ontology Consortium (2000). Gene ontology: tool for the unification of biology. *Nat. Genet.* 25, 25–29.
- Beck, M., Schmidt, A., Malmstroem, J., Claassen, M., Ori, A., Szymborska, A., Herzog, F., Rinner, O., Ellenberg, J., and Aebersold, R. (2011). The quantitative proteome of a human cell line. *Mol. Syst. Biol.* 7, 549.
- Bolender, N., Sickmann, A., Wagner, R., Meisinger, C., and Pfanner, N. (2008). Multiple pathways for sorting mitochondrial precursor proteins. *EMBO Rep.* 9, 42–49.
- Chandra, N.C., Spiro, M.J., and Spiro, R.G. (1998). Identification of a glycoprotein from rat liver mitochondrial inner membrane and demonstration of its origin in the endoplasmic reticulum. *J. Biol. Chem.* 273, 19715–19721.
- Chapman-Smith, A., and Cronan, J.E. (1999). Molecular biology of biotin attachment to proteins. *J. Nutr.* 129, 477S–484S.
- Colombini, M. (1979). A candidate for the permeability pathway of the outer mitochondrial membrane. *Nature* 279, 643–645.
- Csordás, G., Golenár, T., Seifert, E.L., Kamer, K.J., Sancak, Y., Perocchi, F., Moffat, C., Weaver, D., de la Fuente Perez, S., Bogorad, R., et al. (2013). MICU1 controls both the threshold and cooperative activation of the mitochondrial Ca²⁺ uniporter. *Cell Metab.* 17, 976–987.
- Dunkley, T.P.J., Dupree, P., Watson, R.B., and Lilley, K.S. (2004). The use of isotope-coded affinity tags (ICAT) to study organelle proteomes in Arabidopsis thaliana. *Biochem. Soc. Trans.* 32, 520–523.
- Foster, L.J., de Hoog, C.L., Zhang, Y., Zhang, Y., Xie, X., Mootha, V.K., and Mann, M. (2006). A mammalian organelle map by protein correlation profiling. *Cell* 125, 187–199.
- Gebert, N., Ryan, M.T., Pfanner, N., Wiedemann, N., and Stojanovski, D. (2011). Mitochondrial protein import machineries and lipids: a functional connection. *Biochim. Biophys. Acta* 1808, 1002–1011.

- Grynkiewicz, G., Poenie, M., and Tsien, R.Y. (1985). A new generation of Ca²⁺ indicators with greatly improved fluorescence properties. *J. Biol. Chem.* **260**, 3440–3450.
- Hamasaki, M., Furuta, N., Matsuda, A., Nezu, A., Yamamoto, A., Fujita, N., Oomori, H., Noda, T., Haraguchi, T., Hiraoka, Y., et al. (2013). Autophagosomes form at ER-mitochondria contact sites. *Nature* **495**, 389–393.
- Herrmann, J.M., and Riemer, J. (2010). The intermembrane space of mitochondria. *Antioxid. Redox Signal.* **13**, 1341–1358.
- Hoffman, N.E., Chandramoorthy, H.C., Shamugapriya, S., Zhang, X., Rajan, S., Mallilankaraman, K., Gandhirajan, R.K., Vagnozzi, R.J., Ferrer, L.M., Sreekrishnanilayam, K., et al. (2013). MICU1 motifs define mitochondrial calcium uniporter binding and activity. *Cell Rep* **5**, 1576–1588.
- Hong, Y.-C., Leary, S.C., Cobine, P.A., Young, F.B.J., George, G.N., Shoubridge, E.A., and Winge, D.R. (2005). Human Sco1 and Sco2 function as copper-binding proteins. *J. Biol. Chem.* **280**, 34113–34122.
- Itakura, E., Kishi-Itakura, C., and Mizushima, N. (2012). The hairpin-type tail-anchored SNARE syntaxin 17 targets to autophagosomes for fusion with endosomes/lysosomes. *Cell* **151**, 1256–1269.
- Kamer, K.J., and Mootha, V.K. (2014). MICU1 and MICU2 play nonredundant roles in the regulation of the mitochondrial calcium uniporter. *EMBO Rep.* **15**, 299–307.
- Kornmann, B., and Walter, P. (2010). ERMES-mediated ER-mitochondria contacts: molecular hubs for the regulation of mitochondrial biology. *J. Cell Sci.* **123**, 1389–1393.
- Mallilankaraman, K., Doonan, P., Cárdenas, C., Chandramoorthy, H.C., Müller, M., Miller, R., Hoffman, N.E., Gandhirajan, R.K., Molgó, J., Birnbaum, M.J., et al. (2012). MICU1 is an essential gatekeeper for MCU-mediated mitochondrial Ca²⁺ uptake that regulates cell survival. *Cell* **151**, 630–644.
- Martell, J.D., Deerinck, T.J., Sancak, Y., Poulos, T.L., Mootha, V.K., Sosinsky, G.E., Ellisman, M.H., and Ting, A.Y. (2012). Engineered ascorbate peroxidase as a genetically encoded reporter for electron microscopy. *Nat. Biotechnol.* **30**, 1143–1148.
- Mehta, S., and Zhang, J. (2011). Reporting from the field: genetically encoded fluorescent reporters uncover signaling dynamics in living biological systems. *Annu. Rev. Biochem.* **80**, 375–401.
- Neupert, W., and Herrmann, J.M. (2007). Translocation of proteins into mitochondria. *Annu. Rev. Biochem.* **76**, 723–749.
- Ong, S.-E., Blagoev, B., Kratchmarova, I., Kristensen, D.B., Steen, H., Pandey, A., and Mann, M. (2002). Stable isotope labeling by amino acids in cell culture, SILAC, as a simple and accurate approach to expression proteomics. *Mol. Cell. Proteomics* **1**, 376–386.
- Pagliarini, D.J., Calvo, S.E., Chang, B., Sheth, S.A., Vafai, S.B., Ong, S.-E., Walford, G.A., Sugiana, C., Boneh, A., Chen, W.K., et al. (2008). A mitochondrial protein compendium elucidates complex I disease biology. *Cell* **134**, 112–123.
- Perocchi, F., Gohil, V.M., Girgis, H.S., Bao, X.R., McCombs, J.E., Palmer, A.E., and Mootha, V.K. (2010). MICU1 encodes a mitochondrial EF hand protein required for Ca²⁺ uptake. *Nature* **467**, 291–296.
- Polianskyte, Z., Peitsaro, N., Dapkunas, A., Liobikas, J., Soliymani, R., Lalowski, M., Speer, O., Seitsonen, J., Butcher, S., Cereghetti, G.M., et al. (2009). LACTB is a filament-forming protein localized in mitochondria. *Proc. Natl. Acad. Sci. USA* **106**, 18960–18965.
- Rhee, H.-W., Zou, P., Udeshi, N.D., Martell, J.D., Mootha, V.K., Carr, S.A., and Ting, A.Y. (2013). Proteomic mapping of mitochondria in living cells via spatially restricted enzymatic tagging. *Science* **339**, 1328–1331.
- Samali, A., Cai, J., Zhivotovsky, B., Jones, D.P., and Orrenius, S. (1999). Presence of a pre-apoptotic complex of pro-caspase-3, Hsp60 and Hsp10 in the mitochondrial fraction of jurkat cells. *EMBO J.* **18**, 2040–2048.
- Sancak, Y., Markhard, A.L., Kitami, T., Kovács-Bogdán, E., Kamer, K.J., Udeshi, N.D., Carr, S.A., Chaudhuri, D., Clapham, D.E., Li, A.A., et al. (2013). EMRE is an essential component of the mitochondrial calcium uniporter complex. *Science* **342**, 1379–1382.
- Sultan, M., Schulz, M.H., Richard, H., Magen, A., Klingenhoff, A., Scherf, M., Seifert, M., Borodina, T., Soldatov, A., Parkhomchuk, D., et al. (2008). A global view of gene activity and alternative splicing by deep sequencing of the human transcriptome. *Science* **321**, 956–960.
- Tamai, S., Iida, H., Yokota, S., Sayano, T., Kiguchiya, S., Ishihara, N., Hayashi, J.-I., Mihara, K., and Oka, T. (2008). Characterization of the mitochondrial protein LETM1, which maintains the mitochondrial tubular shapes and interacts with the AAA-ATPase BCS1L. *J. Cell Sci.* **121**, 2588–2600.
- Ting, A.Y., Kain, K.H., Klemke, R.L., and Tsien, R.Y. (2001). Genetically encoded fluorescent reporters of protein tyrosine kinase activities in living cells. *Proc. Natl. Acad. Sci. USA* **98**, 15003–15008.
- Vögtle, F.-N., Burkhart, J.M., Rao, S., Gerbeth, C., Hinrichs, J., Martinou, J.-C., Chacinska, A., Sickmann, A., Zahedi, R.P., and Meisinger, C. (2012). Intermembrane space proteome of yeast mitochondria. *Mol. Cell. Proteomics* **11**, 1840–1852.