



**A**

	A	B	C	D	E	F	G	H	I	J	K	L
A	0	0.001	0.001	0.003	0.003	0.003	0.333	0.334	0.335	0.335	0.336	0.336
B	11.5	0	0.000	0.002	0.002	0.002	0.332	0.333	0.334	0.334	0.335	0.336
C	11.5	3.0	0	0.002	0.002	0.002	0.332	0.333	0.334	0.334	0.335	0.336
D	27.5	22.5	22.5	0	0.000	0.000	0.331	0.331	0.332	0.332	0.333	0.334
E	27.5	18.0	22.5	1.0	0	0.000	0.331	0.331	0.332	0.332	0.333	0.334
F	27.5	22.5	18.0	3.0	3.0	0	0.331	0.331	0.332	0.332	0.333	0.334
G	50.0	42.0	42.0	32.0	32.0	32.0	0	0.000	0.001	0.002	0.003	0.003
H	53.5	47.0	47.0	34.5	34.5	36.0	6.0	0	0.001	0.001	0.002	0.003
I	60.0	53.5	53.5	38.0	38.0	38.0	14.0	9.0	0	0.001	0.001	0.002
J	62.0	57.5	57.5	42.0	42.0	42.0	18.0	14.0	6.0	0	0.001	0.002
K	64.0	60.0	60.0	47.0	47.0	47.0	25.0	18.0	9.0	6.0	0	0.001
L	66.0	64.0	64.0	53.5	53.5	53.5	30.0	27.5	18.0	14.0	9.0	0



©Figure 16.6 McGarvey et al. 2017. *In: Hauer and Lamberti (eds.) Methods in Stream Ecology*. Elsevier.