



Landscape factors and hydrology influence mercury concentrations in wading birds breeding in the Florida Everglades, USA



Garth Herring ^{a,*}, Collin A. Eagles-Smith ^a, Joshua T. Ackerman ^b, Dale E. Gawlik ^c, James M. Beerens ^c

^a U.S. Geological Survey, Forest and Rangeland Ecosystem Science Center, 3200 SW Jefferson Way, Corvallis, OR 97331, USA

^b U.S. Geological Survey, Western Ecological Research Center, Dixon Field Station, 800 Business Park Drive, Suite D, Dixon, CA 95620, USA

^c Department of Biological Sciences, Florida Atlantic University, 777 Glades Rd, Boca Raton, FL 33440, USA

HIGHLIGHTS

- Hydrologic variables may influence Hg exposure in wading birds.
- Hg in adult great egrets was most influenced by region.
- Adult white ibis Hg was most influenced the number of days a site was dry.
- Hg in egret chicks was most influenced by date.
- Hg in ibis chicks was most influenced by chick age, region, and water recession rate.

ARTICLE INFO

Article history:

Received 7 January 2013

Received in revised form 10 April 2013

Accepted 11 April 2013

Available online 23 May 2013

Keywords:

Great egret
Hydrology
Methylmercury
Everglades
Wetlands
White ibis

ABSTRACT

The hydrology of wetland ecosystems is a key driver of both mercury (Hg) methylation and waterbird foraging ecology, and hence may play a fundamental role in waterbird exposure and risk to Hg contamination. However, few studies have investigated hydrological factors that influence waterbird Hg exposure. We examined how several landscape-level hydrological variables influenced Hg concentrations in great egret and white ibis adults and chicks in the Florida Everglades. The great egret is a visual “exploiter” species that tolerates lower prey densities and is less sensitive to hydrological conditions than is the white ibis, which is a tactile “searcher” species that pursues higher prey densities in shallow water. Mercury concentrations in adult great egrets were most influenced by the spatial region that they occupied in the Everglades (higher in the southern region); whereas the number of days a site was dry during the previous dry season was the most important factor influencing Hg concentrations in adult ibis (Hg concentrations increased with the number of days dry). In contrast, Hg concentrations in egret chicks were most influenced by calendar date (increasing with date), whereas Hg concentrations in ibis chicks were most influenced by chick age, region, and water recession rate (Hg concentrations decreased with age, were higher in the southern regions, and increased with positive water recession rates). Our results indicate that both recent (preceding two weeks) hydrological conditions, and those of the prior year, influence Hg concentrations in wading birds. Further, these results suggest that Hg exposure in wading birds is driven by complex relationships between wading bird behavior and life stage, landscape hydrologic patterns, and biogeochemical processes.

Published by Elsevier B.V.

1. Introduction

A greater understanding of the global distribution and atmospheric deposition of mercury (Hg) has led to heightened concern for wildlife exposure, because as impairment thresholds become refined and better quantified, there is increasing evidence that environmentally relevant Hg exposure can result in impaired reproduction and other toxicological endpoints such as endocrine disruption (Frederick and Jayasena, 2010; Jayasena et al., 2011). Waterbirds are particularly prone to elevated Hg exposure and associated toxicity in wetland ecosystems (Frederick et al.,

2002; Eagles-Smith et al., 2009; Ackerman et al., 2011) where they often are top predators, and biogeochemical conditions facilitate the conversion of inorganic Hg to methylmercury (MeHg; Marvin-DiPasquale et al., 2003; Hall et al., 2008), the most toxic and bioavailable form. Wetlands with substantial hydrologic fluctuations can potentially enhance both the methylation and availability of MeHg (Cleckner et al., 1998; Gustin et al., 2006). Consequently, waterbirds that rely on ephemeral hydrology for enhanced foraging opportunities can be exposed to elevated MeHg through food web biomagnification (Sundlof et al., 1994; Frederick et al., 2002; Custer et al., 2008).

Shallow wetland ecosystems with regularly fluctuating hydrologic conditions and elevated loading of reactive inorganic Hg also present an ideal situation to study landscape hydrologic factors that may

* Corresponding author. Tel.: +1 541 750 0952.

E-mail address: gherring@usgs.gov (G. Herring).

promote the exposure and uptake of MeHg by upper trophic level waterbirds. The Florida Everglades is particularly well suited for this type of study because it is a very large (8713 km²) pulsed hydrologic system (wet and dry seasons; Obeysekerera et al., 1999), and over 60% of the landscape dries out each year (USGS, 2006). The Everglades has a well-documented history of elevated atmospheric Hg deposition and conversion of inorganic Hg to MeHg on a landscape scale (Osborne et al., 2011), and supports extensive populations of breeding wading birds which have previously been shown to have elevated Hg concentrations (Sundlof et al., 1994; Rumbold et al., 2001; Frederick et al., 2002; Herring et al., 2009). Importantly, large areas of the Everglades contain elevated dissolved sulfate concentrations (20–50 mg/l) that exceed background levels (<1 mg/l) (Orem et al. 2011). Sulfate in the Everglades originates from the Everglades Agricultural Area and is associated with both agricultural practices and from the sulfur released by oxidation of organic soils (Orem et al. 2011). Sulfate plays a critical role in the production of MeHg, via microbial sulfate reduction, resulting in increased bioavailability and biomagnification of MeHg (Gilmour et al., 2007; Orem et al., 2011). Further, periphyton which serves as the base of the Everglades food web (Cleckner et al. 1998), has relatively high MeHg concentrations (Liu et al. 2008), and enhances the bioavailability of Hg in Everglades species (Cleckner et al., 1998; Molina et al., 2010). Thus, the hydrological pulsed nature of the Everglades, elevated atmospheric wet deposition of Hg (Mercury Deposition Network, 2012), periphyton based food web, and high concentrations of sulfate result in considerable MeHg production and subsequent availability within Everglades food chains.

Wading bird reproduction in the freshwater region of the Everglades occurs during the dry season (Dec–June), when water levels recede across the landscape, increasing aquatic prey density in shallow depressions (Kushlan, 1981; Gawlik, 2002). As such, hydrologic patterns are the dominant driver of wading bird foraging locations (Bancroft et al., 2002; Russell et al., 2002; Beerens et al., 2011), reproductive success (Kushlan, 1987; Frederick and Collopy, 1989; Herring et al., 2010), and health (Herring et al., 2011; Herring and Gawlik, 2012). Importantly, hydrologic factors associated with wading bird foraging patterns also may be associated with Hg methylation. The influence of hydrologic factors on waterbird MeHg exposure is poorly understood, but has important ramifications for water management activities in wetland ecosystems.

We evaluated MeHg exposure in two species of wading birds in relation to landscape-level hydrologic factors that are known to influence either MeHg production or wading bird foraging ecology. The great egret (*Ardea alba*; hereafter egret) and white ibis (*Eudocimus albus*; hereafter ibis) breed extensively throughout the Everglades and serve as long-term indicator species for Hg risk to the Everglades ecosystem (see Sundlof et al., 1994; Rumbold et al., 2001; Frederick et al., 2002; Heath and Frederick, 2005; Herring et al., 2009). Accordingly, they serve as important indicator species for the Everglades ecosystem. Although egrets and ibises nest and breed and feed sympatrically (that is coexist), they differ both in their foraging strategies (Gawlik, 2002; Beerens et al., 2011) and diets (Smith, 1997; Dorn et al., 2011). Ibises are “searchers” who target ephemeral high-quality patches of biomass, then abandon them once prey biomass declines, whereas egrets are “exploiters” that continue foraging at sites even when prey densities are low (Gawlik, 2002). Suitable foraging conditions for ibises also span a much narrower range of water depths than egrets (Beerens et al., 2011). Further, ibises usually consume crayfish (*Procambarus* spp.) but can switch to consuming fish when water recession rates and depths produce high fish densities (Kushlan, 1979; Dorn et al., 2011). In contrast, egrets select larger prey items than ibis (generally fish) regardless of habitat conditions (Smith, 1997). We hypothesized that differences in prey resources and habitat conditions (e.g., depths) would result in differential Hg exposure between these two species. Further, there may be differences in prey delivery to chicks of both species associated with chick size and semi-altricial (down covered at birth, nest bound, and fed by parents) life history of both species (Heath et al., 2009; McCrimmon et al., 2011), as well as associated differences in diets delivered to chicks

under varying hydrological conditions and landscape level prey densities (Dorn et al., 2011). Our objectives in this study were to assess the relative importance of factors associated with foraging habitat conditions, the production of MeHg, and large scale regional MeHg patterns on Hg concentrations in great egrets and white ibises, and to determine whether the drivers of Hg exposure are similar across adult and chick life stages.

2. Materials and methods

2.1. Study area

The Florida Everglades is a large, subtropical and oligotrophic wetland in southern Florida with pronounced annual wet and dry seasons (Obeysekerera et al., 1999). The dry season, when most wading birds breed, typically spans from November through May. The Everglades ecosystem has been highly altered over the past 60 years, with more than half of its spatial extent lost to agriculture and urban development (Ogden et al., 2003). As a result, the hydrologic patterns of remnant areas are greatly altered, with changes to the timing, magnitude and predictability of both the hydrology and seasonal prey availability events that are critical to wading bird reproduction (Ogden, 1994). Although Hg concentrations in the Everglades wildlife have generally declined over the past decade, some areas still pose reproductive risks to piscivorous birds (Rumbold et al., 2008).

2.2. Bird capture and sample collection

2.2.1. Adult sampling

Adult egrets and ibises were captured during the pre-breeding season (Jan 10 to Mar 23 in 2006 and 2007) using either a netgun or modified flip trap (Herring et al., 2008a) and decoys (Crozier and Gawlik, 2003; Heath and Frederick, 2003). Birds were captured at 21 sites in the northern Everglades, within Water Conservation Areas (WCA) 2A, 3A, and the Arthur R. Marshall Loxahatchee National Wildlife Refuge (Lox; Fig. 1). We attached radio transmitters (16 g) using a figure-8 harness to all birds to quantify foraging movement patterns throughout the breeding season and to estimate hydrologic conditions at foraging sites and within foraging ranges of breeding colonies.

We used blood as our matrix to measure Hg because it reflects recent exposure (Bearhop et al., 2000), whereas feather Hg concentrations are reflective of an equilibrium of both Hg exposure and body burden depuration only during the period of feather growth. Therefore, we could not accurately link feather Hg concentrations with recent foraging. From each bird, we collected up to 2 ml of blood from the brachial vein using a 27.5-gauge needle. Blood samples were stored in heparinized cryovials and placed on ice until transported to the lab. The blood volume available for collection while following American Veterinary Medical Association guidelines was limited due to the need to measure other physiological parameters (see Herring and Gawlik, 2012), therefore we measured the red blood cell (RBC) fraction for total Hg concentrations because most of the Hg in blood is in the RBC fraction, and RBC THg concentrations are correlated with the more commonly measured whole blood matrix (Chen et al., 2011; Lieske et al., 2011). Plasma and red blood cells were separated in the lab via centrifugation (15 min, 5000 rpm), and each fraction was frozen at –20 °C for later analysis. All adult birds were later sexed using DNA analysis (Zoogen Services Inc., Davis, CA; Herring et al., 2008b).

2.2.2. Chick sampling

We monitored nests within egret and ibis colonies that were associated both with our radio-tagged adults and easily accessible (Herring et al., 2010; Beerens et al., 2011; Fig. 1). Those nests (egrets = 59, ibises = 70) were revisited approximately every 7 days, and the order of chicks was determined by the hatching sequence noted during visits. Upon arrival at each monitored nest, we immediately removed the nestlings by hand and held them in either a cloth bag or large plastic container to keep




Fig. 1. Location of study area indicating adult wading bird capture sites and wading bird nesting colonies sampled during 2006 and 2007. The primary Water Management Areas (WCAs), Arthur R. Marshall Loxahatchee National Wildlife Refuge (Lox), and Everglades National Park (ENP) are indicated.

them calm and minimize movement during subsequent sampling and measurements. Blood collection and storage followed those methods described for adults above.

2.3. Hydrological conditions

We used the Everglades Depth Estimation Network (EDEN; USGS, 2006) to estimate water depths, water level recession rates, and the number of days sites were dry during the previous dry season (days dry) at both foraging sites where adults were captured and their associated breeding colonies where chicks were sampled. The EDEN uses a network of water level gauges to produce a water surface model that, when combined with a ground elevation model, provides an estimate of water depth for the entire freshwater portion of the Greater Everglades (Liu et al., 2009). EDEN calculates water level depths in $400\text{ m} \times 400\text{ m}$ grid cells at daily time steps accounting for evapotranspiration, rainfall, and sheet flow. The estimated water depths are accurate to within 5 cm (Liu et al., 2009).

We estimated water recession rates, water depths, and days dry at foraging sites by first delineating the EDEN grid cells located within a 3-km radius of each bird capture site using ArcGIS 9.1 (ESRI, Redlands, CA). The 3-km radius was used because, in a prior study, radio-tagged egrets and ibises primarily foraged within 3 km of the capture site for up to 2 weeks prior to capture (Herring and Gawlik, 2012). We then extracted daily water depth values for the 14 days preceding each bird's capture for all cells within the foraging area, and used the mean depth and change-in-depth over each period as the measures of water depth and recession rate, respectively. Positive recession rates indicate decreasing water levels. We estimated days dry by counting the total number of days within the previous dry season that a site remained dry (i.e., depth < 0 cm). Surface water could still be present when estimated cell depth < 0 cm; however, the 5-cm error in EDEN depths suggests that a majority of the ground surface would be exposed in a given $400\text{ m} \times 400\text{ m}$ grid cell.

To estimate recession rates, water depths, and days dry associated with chicks sampled later in the breeding season we first used ArcGIS

9.1 (ESRI, Redlands, CA) to establish fixed radius buffers (hereafter foraging range) around colonies based on mean distances flown + 1 SE by the radio tagged adult egrets and ibises described above. We defined foraging range as the average distance that radio tagged individuals flew from breeding colonies to a foraging site throughout the breeding season. Egrets and ibises are central place foragers, and the use of foraging ranges was supported by our telemetry data. Estimates of foraging distances from colonies were obtained by locating the adult radio-tagged egrets ($n = 76$) and ibises ($n = 125$) via aerial surveys during 2006–2007 using a left–right control box and two side view directional yagi antennas. A random subset (20–30) of all radio-tagged adult egrets and ibises were aerially tracked each day during the morning 3–4 days per week throughout the breeding season to foraging sites. After each flight all located birds were removed from the tracking list, and after two successive aerial surveys all birds were added back into the tracking list. This approach resulted in individual birds being located no more than two times per week, with a minimum of three days between locations to ensure independence. Locations of foraging birds derived from radios were verified visually. The mean distance that adults flew from their nesting colony in 2006 and 2007, respectively, was $12.5 \text{ km} \pm 0.4 \text{ SE}$ ($n = 227$) and $7.4 \pm 0.5 \text{ SE}$ ($n = 178$) for ibises, and $8.1 \text{ km} \pm 0.7 \text{ SE}$ ($n = 60$) and $4.6 \text{ km} \pm 0.3 \text{ SE}$ ($n = 163$) for egrets (Beerens, 2008). We extracted the water depth from each EDEN grid cell within the foraging ranges for specific colonies by year and species. For each day that we sampled nestling blood, we calculated water recession rate and water depth as the mean value for the preceding 2-week period.

2.4. Mercury determination

We analyzed red blood cells for total mercury (THg) because the majority of Hg in blood is in the red blood cell fraction (Myers and Davidson, 1998; Day et al., 2007), and almost all (>95%) of the mercury in avian blood is in the methylmercury form (Thompson and Furness, 1989; Fournier et al., 2002). To minimize any potential variability in THg concentrations associated with differential moisture loss during long-term storage of samples, red blood cell fractions were dried (48 h at 50 °C) to a constant weight before THg analysis. We measured up to 0.05 g of dried blood into a quartz sample vessel (weighed to the nearest 0.00001 g; Mettler Toledo model XS105, Mettler Toledo, Columbus, Ohio, USA). Following U.S. Environmental Protection Agency method 7473 (US EPA, 2000), we determined THg concentrations in each red blood cell sample at the USGS Contaminants Ecology Lab in Corvallis, Oregon on a Milestone DMA-80 direct mercury analyzer (Milestone, Monroe, Connecticut, USA). Analytical equipment was calibrated using certified standard solutions prior to analysis, and accuracy and precision were evaluated within each analytical batch through the inclusion of certified reference materials (either dogfish muscle tissue [DORM-3] or dogfish liver [DOLT-4] by the National Research Council of Canada, Ottawa, Canada), calibration verifications (liquid standards), duplicates, and blanks. Batches consisted of 30 total samples, two of those with duplicates for a total of 32 samples per batch. The remaining 8 places in the DMA sample tray were used for quality assurance and quality control. Recoveries averaged $102.3 \pm 0.07\%$ ($n = 30$) and $96.4 \pm 0.01\%$ ($n = 44$) for certified reference materials and calibration checks, respectively. Absolute relative percent difference for all duplicates averaged $3.1\% \pm 0.12\%$ ($n = 27$).

2.5. Statistical analyses

We used an information-theoretic approach (Burnham and Anderson, 2002) and linear mixed models in R (LME4; R Development Core Team, 2011) to determine the variables that most influenced THg concentrations in adults and chicks of both egrets and ibises. We specified the maximum likelihood estimation method in our linear models. We built and ranked separate competing candidate models for each species and life stage (adult and chick), in order to understand

how their individual Hg concentrations responded to differing habitat conditions (i.e., hydrology and date within years). We also included an intercept only (null) model and global model for each species and life stage. Mercury concentrations were natural log transformed to improve normality of residuals. Where appropriate we included linear, quadratic, and cubic forms of variables (e.g., age). Mercury values are presented as red blood cell THg ($\mu\text{g/g dw} \pm$ standard error [SE]).

We assessed the influence of variables potentially affecting Hg concentrations in two stages; first we assessed those variables associated with just adults, and in the second stage we examined variables potentially associated with Hg concentrations in chicks. Variables were included that could result in egrets and ibises being exposed to differing levels of Hg through their foraging habitat preferences and habitat conditions (date, water recession rate, water depth; Gawlik, 2002; Beerens et al., 2011), prey selection behavior (year, sex, age of chicks; Smith, 1997; Loftus, 2000; Dorn et al., 2011), landscape factor related to the production of MeHg (days dry, region; Krabbenhoft and Fink, 2001; Axelrad et al., 2007; Gilmour et al., 2007), factors related to the dilution of Hg associated with chick growth (age; Ackerman et al., 2011), and large scale regional MeHg patterns (region; Herring et al., 2009). Prior to the model selection analyses we examined correlations between all variables to detect any autocorrelation issues. For adults there were no autocorrelation issues (all $r^2 \leq 0.30$). However, for chicks we observed a positive correlation between age and date ($r^2 = 0.40$ – 0.69) because nesting was moderately synchronous during our study. To determine which variable to include in our candidate models we ran a preliminary model sets for each species that included all variables but did not allow age and date to be in the same model, we compared the most parsimonious model (included date for egrets and age for ibises) with the first model where age or date was replaced by the reciprocal. For both egrets ($\Delta\text{AIC}_c = 12.76$) and ibises ($\Delta\text{AIC}_c = 18.03$), there was strong support for retaining either date or age respectively, but not both variables in their respective model sets. For subsequent egret models we included only the date variables, whereas for ibis chicks we included the age variable only. For adults we included seven model variables: year, region, water recession rate, water depth, date, days dry, and sex. In the case of chicks we included seven model variables: year, region, water recession rate, water depth, days dry, hatch order and either date (egrets) or age (ibises). We included the following random effects; colony location (colony) to account for sampling different numbers of nests within colonies, chick identification number to account for the repeated sampling of chicks, and nest identification to account for the non-independence of sampling multiple chicks within a nest.

We used Akaike's information criterion adjusted for small samples sizes (AIC_c) and considered the model with the lowest AIC_c value to be the most parsimonious (Burnham and Anderson, 2002). We determined the relative ranking of each model by subtracting each candidate model's AIC_c value from the best model (ΔAIC_c). We ran all possible models and considered candidate models with $\Delta\text{AIC}_c \leq 2.0$ to be equally plausible and models with $\Delta\text{AIC}_c > 4.0$ to have less support. We present all candidate models with $\Delta\text{AIC}_c \leq 4.0$ and the null model in tables. We calculated Akaike weights (w_i) to assess the weight of evidence for candidate models (Burnham and Anderson, 2002). We also calculated cumulative variable weights by summing Akaike weights across all models that included the variable to assess the relative importance of each variable. Where multiple forms of a variable (age) were included (linear, quadratic, and cubic) we present only the variable with the highest relative variable weight. In this case the only way for the cubic effect of age to be included is with both the linear and quadratic effects. We calculated model-averaged beta coefficient estimates from all candidate models (Burnham and Anderson, 2002) and present them with 95% confidence intervals.

3. Results

During the 2006 and 2007 pre-breeding seasons we captured and collected blood from 187 adults (68 egrets; 2006 = 22, 2007 = 46, and 119

ibises; 2006 = 54, 2007 = 65). The geometric mean red blood cell THg concentration (µg/g dw) averaged across dates, regions, sex, and years was 4.46 ± 0.40 (SE) in egret adults and 2.44 ± 0.12 in ibis adults. During the 2006 and 2007 nesting seasons we collected blood samples from 254 chicks (169 egret; 2006 = 28, 2007 = 147, and 85 ibis; 2006 = 30, 2007 = 55). The geometric mean red blood cell THg concentration (µg/g dw) averaged across age, dates, hatch order, regions, and years was 4.14 ± 0.16 in egret chicks, and 0.58 ± 0.07 in ibis chicks.

3.1. Adult mercury

3.1.1. Great egret

The most parsimonious model explaining adult egret THg contained the variables for region and recession rate with an Akaike weight (w_i) of 0.14 (Table 1). Two additional models containing region, recession rate, and water depth, and region, recession rate, and sex also had reasonable although the log-likelihood values for these competing models did not improve substantially (<2 ΔAIC_c units, Table 1). Using cumulative variable weights to assess variable importance, we found that models containing region had a high level of support with a combined AIC_c weight of 0.91, followed by moderate support for recession rate (0.76), with little evidence for effects of sex (0.31), water depth (0.30), year (0.27), date (0.26), or days dry (0.24). The model-averaged estimate for adult egret THg concentrations by region was 416% higher at WCA 3A than Lox, 61% higher at WCA 3A than WCA 2A, and 221% higher in WCA 2A than Lox (Fig. 2). Although recession rate had a moderately strong variable weight, the model averaged beta coefficient estimate

had 95% confidence intervals that overlapped zero, indicating recession rate ($\beta = 0.410 \pm 0.421$) had a smaller effect on adult egret Hg concentrations than did region ($\beta = 1.733 \pm 1.131$).

3.1.2. White ibis

The most parsimonious model explaining adult ibis Hg concentrations contained the variables for days dry, date, and water depth with an Akaike weight of 0.15 (Table 1). Two additional models containing days dry, date, water depth, and year and days dry, date, water depth, and sex also had reasonable support although the log-likelihood values for these competing models did not improve substantially (<2 ΔAIC_c units, Table 1). Using cumulative variable weights to assess variable importance, we found that models containing days dry had a high level of support with a combined AIC_c weight of 0.95, followed by date (0.73), and water depth (0.64), with little evidence for effects of year (0.40), sex (0.28), recession rate (0.26), or region (0.20). Model-averaged estimates for the effect of days dry on adult ibis Hg concentrations indicated an increase in Hg concentrations of 119% across an increasing numbers of days a site was dry prior to the proceeding rainy season (range = 4–109 days, $\beta = 0.007 \pm 0.005$; Fig. 3).

Although date and water depth had moderate variable weights, the model averaged beta coefficient estimates for these variables had 95% confidence intervals that overlapped zero indicating that both date ($\beta = -0.007 \pm 0.008$) and water depth ($\beta = 0.010 \pm 0.013$) had a much smaller effect on adult white ibis Hg concentrations than did days dry ($\beta = 0.007 \pm 0.005$).

Table 1

Ranking of candidate models describing variables influencing Hg concentrations in adult great egrets and white ibises in the Florida Everglades.^a Models are ranked by differences in Akaike's information criterion. Only candidate models within ΔAIC_c ≤ 4.0 and the null model are presented.

Model structure	K ^b	-2LogL	AIC _c ^c	ΔAIC _c ^d	w _i ^e	Evidence ratio ^f
<i>Great egrets (n = 62)</i>						
Region + recession rate	5	119.45	130.52	0.00	0.14	1.00
Region + recession rate + water depth	6	118.32	131.84	1.32	0.07	1.94
Region + recession rate + sex	6	118.69	132.22	1.69	0.06	2.33
Region + recession rate + days dry	6	119.17	132.70	2.18	0.05	2.97
Region + recession rate + date	6	119.26	132.79	2.26	0.05	3.10
Region + recession rate + year	6	119.27	132.80	2.27	0.05	3.12
Region	4	124.61	133.31	2.79	0.04	4.03
Region + recession rate + water depth + sex	7	117.53	133.61	3.09	0.03	4.68
Region + sex	5	122.95	134.02	3.50	0.02	5.76
Region + recession rate + water depth + year	7	118.25	134.33	3.80	0.02	6.70
Region + recession rate + water depth + days dry	7	118.30	134.37	3.85	0.02	6.85
Region + recession rate + water depth + date	7	118.32	134.39	3.87	0.02	6.92
Intercept	2	135.24	139.44	9.84	0.00	136.89
<i>White ibises (n = 105)</i>						
Days dry + date + water depth	5	170.75	181.36	0.00	0.15	1.00
Days dry + date + water depth + year	6	169.88	182.74	1.38	0.07	1.99
Days dry + date + water depth + sex	6	170.08	182.94	1.58	0.07	2.21
Days dry + date + water depth + recession rate	6	170.70	183.56	2.20	0.05	3.01
Days dry + date + year	5	173.09	183.70	2.34	0.05	3.23
Days dry + date	4	175.37	183.77	2.42	0.04	3.35
Days dry	3	177.61	183.85	2.50	0.04	3.49
Days dry + water depth + year + region	7	169.27	184.43	3.07	0.03	4.65
Days dry + date + year + sex	7	169.49	184.65	3.29	0.03	5.18
Days dry + date + water depth + year + recession rate	7	169.86	185.01	3.66	0.02	6.23
Days dry + year	4	176.63	185.03	3.67	0.02	6.28
Days dry + water depth	4	176.81	185.21	3.85	0.02	6.87
Days dry + date + water depth + sex + recession rate	7	170.06	185.21	3.86	0.02	6.89
Days dry + date + water depth + region	7	170.20	185.36	4.00	0.02	7.39
Intercept	2	196.13	200.25	18.89	0.00	7.82E ± 06

^a Adults were sampled during the pre-breeding stage across Water Conservation Areas 1–3 during 2006–2007. Variables considered included: region, date captured, number of days a site was dry the previous dry season (days dry), water depth, recession rate, year, and sex.

^b Number of estimated parameters in the model including the intercept and variance.

^c Second-order Akaike's information criterium (AIC_c).

^d The difference in the value between AIC_c of the current model and the value of the most parsimonious model.

^e Likelihood of the model given the data, relative to candidate models.

^f The weight of the evidence that the top model is better than the selected model, given the candidate model set.




Fig. 2. Model averaged least square mean red blood cell Hg ($\mu\text{g/g dw}$) \pm SE concentrations in great egret and white ibis adults and chicks in the Arthur R. Marshall Loxahatchee National Wildlife Refuge (Lox), and Water Conservation Areas 2A and 3A (WCA 2A, WCA 3A) during 2006 and 2007. No ibis chicks were sampled in WCA 2A during the study.

3.2. Chick mercury

3.2.1. Great egrets

The most parsimonious model explaining egret chick Hg concentrations contained the variables for date, days dry, and year with an Akaike weight of 0.09 (Table 1). A number of other models provided a reasonably good fit to the data although the log-likelihood values for these competing




Fig. 3. Relationship between partial residuals of red blood cell Hg value ($\mu\text{g/g dw}$) and days dry (number of days a site was dry during the previous dry season) for adult white ibises in the Florida Everglades, after accounting for the other predictor variables in the best models (see Table 1). THg concentrations increased by 119% across the range of day sites were dry (cumulative parameter weight = 0.95, $\beta = 0.007 \pm 0.005$).

models did not improve substantially ($<2 \Delta\text{AIC}_c$ units, Table 1). Using cumulative variable weights to assess variable importance, we found that models containing date had a high level of support with a combined AIC_c weight of 1.0, with moderate support for days dry (0.53), and little support for year (0.43), recession rate (0.33), water depth (0.30), hatch order (0.19), or region (0.10). The model averaged beta coefficient estimate for date on egret chick Hg concentrations ($\beta = 0.026 \pm 0.005$) indicated an increase in Hg concentrations of 326% across the observed range of dates (range = April 14–June 8; Fig. 4). Although days dry had a moderate variable weight the model averaged beta coefficient estimate for this variable had 95% confidence intervals that slightly overlapped zero, indicating that days dry ($\beta = 0.003 \pm 0.005$) had much smaller effect on egret chick Hg concentrations than did date.

3.2.2. White ibises

The most parsimonious model explaining ibis chick Hg contained the variables for age, age², age³, region, and recession rate with an Akaike weight of 0.36 (Table 2). One other candidate model provided a reasonably good fit to the data although the log-likelihood values for these competing models did not improve substantially ($<2 \Delta\text{AIC}_c$ units, Table 2). Using cumulative variable weights to assess variable importance, we found a high degree of support for models containing age³ (AIC_c weight of 0.99), followed by region (0.91), recession rate (0.84), and little support for water depth (0.37), days dry (0.19), hatch order (0.05), or year (0.01). Model-averaged estimates for the effect of age on white ibis chick Hg concentrations indicated a two-fold decrease in Hg concentrations across the observed range of ages (range = 3–30 days, $\beta = -0.001 \pm 0.000$; Fig. 5A). Ibis chick Hg concentrations were 126% higher in WCA 3A than Lox (Fig. 2), and increased by 162% as water recession rates increased (range = -2.29 – 1.58 cm/d, $\beta = 0.248 \pm 0.201$; Fig. 5B).

4. Discussion

4.1. Factors influencing adult egret and ibis mercury concentrations

These results highlight the complexities of Hg cycling through food webs and the interaction between biogeochemical processes, landscape processes, and foraging responses (Fig. 6). Although there was substantial overlap in the sampling sites for each species, the factors influencing their Hg concentrations were different and depended on both foraging strategies and factors likely associated with landscape production of MeHg. In the case of egrets, region was found to be the most important variable influencing their Hg concentrations. Similar to previous studies of wading bird Hg exposure in the Everglades (see Herring et al., 2009), we found that THg concentrations were higher in WCA 3A than WCA




Fig. 4. Relationship between partial residuals of red blood cell Hg value ($\mu\text{g/g dw}$) and the date great egret chicks were sampled in the Florida Everglades, after accounting for the other predictor variables in the best models (see Table 2). THg concentrations increased by 326% across the range of dates (cumulative parameter weight = 1.0, $\beta = 0.026 \pm 0.005$).

Table 2

Ranking of candidate models describing variables influencing Hg concentrations in great egret and white ibis chicks in the Florida Everglades.^a Models are ranked by differences in Akaike's information criterion. Only candidate models within $\Delta AIC_c \leq 4.0$ and the null model are presented.

Model structure	K^b	-2LogL	AIC_c^c	ΔAIC_c^d	w_i^e	Evidence ratio ^f
<i>Great egrets (n = 169)</i>						
Date + days dry + year	7	229.01	243.71	0.00	0.09	1.00
Date	5	233.57	243.94	0.23	0.08	1.12
Date + days dry	6	231.55	244.07	0.37	0.08	1.20
Date + year	6	232.04	244.56	0.86	0.06	1.53
Date + recession rate	6	232.24	244.76	1.06	0.05	1.70
Date + days dry + year + recession rate	8	228.02	244.92	1.21	0.05	1.83
Date + days dry + water depth	7	230.41	245.11	1.40	0.05	2.02
Date + days dry + recession rate	7	230.54	245.24	1.53	0.04	2.15
Date + year + recession rate	7	230.67	245.36	1.66	0.04	2.29
Date + water depth	6	232.96	245.48	1.78	0.04	2.43
Date + days dry + year + water depth	8	228.95	245.85	2.15	0.03	2.93
Date + recession rate + water depth	7	231.57	246.27	2.56	0.03	3.60
Date + days dry + recession rate + water depth	8	229.42	246.32	2.61	0.03	3.69
Date + hatch order	7	231.75	246.45	2.74	0.02	3.94
Date + days dry + hatch order	8	229.65	246.55	2.84	0.02	4.14
Date + year + water depth	7	232.03	246.73	3.02	0.02	4.53
Date + days dry + year + hatch order	9	227.88	247.02	3.31	0.02	5.23
Date + days dry + year + recession rate + water depth	9	227.97	247.10	3.40	0.02	5.47
Date + recession rate + hatch order	8	230.40	247.30	3.60	0.02	6.03
Date + year + recession rate + water depth	8	230.65	247.55	3.85	0.01	6.84
Date + year + hatch order	8	230.80	247.70	3.99	0.01	7.36
Intercept	4	279.94	288.18	44.48	0.00	4.55E + 09
<i>White ibises (n = 85)</i>						
Age + age ² + age ³ + region + recession rate	9	202.19	222.59	0.00	0.36	1.00
Age + age ² + age ³ + region + recession rate + depth	10	200.37	223.35	0.75	0.25	1.46
Age + age ² + age ³ + region + recession rate + days dry	10	202.10	225.07	2.48	0.10	3.45
Age + age ² + age ³ + region	8	208.42	226.32	3.72	0.06	6.43
Intercept	4	249.96	258.46	35.87	0.00	5.85E ± 09

^a Chicks were sampled 12 breeding colonies across Water Conservation Areas 1–3 during 2006–2007. Variables considered included: region, date sampled, number of days a site was dry the previous dry season (days dry), water depth, recession rate, year, and hatch order.

^b Number of estimated parameters in the model including the intercept and variance.

^c Second-order Akaike's information criterion (AIC_c).

^d The difference in the value between AIC_c of the current model and the value of the most parsimonious model.

^e Likelihood of the model given the data, relative to models in the candidate set.

^f The weight of the evidence that the top model is better than the selected model, given the candidate model set.

2A or Lox in adult egrets. Egrets have the flexibility to exploit sites with deeper water (Gawlik, 2002), and a wider range of prey densities (Herring et al., 2010; Herring and Gawlik, 2012), and subsequently can spend more time in regions of the Everglades whereas ibises cannot. Thus, regional differences in adult egret THg appear to be less influenced by the specific hydrological variables we included in our models than the long-term overarching factor of region (Lox, WCA 2A, WCA 3A). Specifically, MeHg concentrations in soils and biota tend to be lower in areas of decreased wetland disturbance (e.g., Lox) and higher in more eutrophic areas of WCA 2A and WCA 3A (Axelrad et al., 2007). These results may also be exacerbated by sulfate concentrations that are higher in WCA 2A and WCA 3A than Lox (Axelrad et al., 2007), because sulfate has been shown to be a limiting factor in the cycling of mercury in the Everglades (Gilmour et al., 2004; Gilmour et al., 2007; Orem et al., 2011).

Conversely, regardless of region, THg concentrations in adult ibis increased in individuals foraging in sites with greater number of days dry, suggesting that this relationship may be driven by the complex Hg methylation process. Everglades soils that become dry for longer periods of time may have more oxidation of sulfides to sulfate. Moreover, Hg in drier soils is more likely oxidize to reactive Hg(II) forms (Krabbenhoft and Fink, 2001). Together, these processes will facilitate the production of MeHg once soils reflood during the wet season, providing anaerobic microbial communities the requisite electron acceptors and bioavailable inorganic Hg needed for elevated MeHg production (Gilmour et al., 2007; Orem et al., 2011). Aquatic prey species with limited movement during the dry season (e.g., crayfish, small fishes) would then be exposed to spikes in MeHg at these sites when they flood, resulting in very specific Hg exposure patterns in wading birds that target wetland areas with these specific hydrology conditions (i.e., short hydroperiod wetlands

that regularly dry out each year). We hypothesize that Hg continues to increase with the number of days sites are dry for some period of time (up to 110 days in our study) because when the average depth of a 400 m × 400 m EDEN cell is 0 cm, the higher elevation portions of that cell have likely been dry several weeks (oxidation of sulfides may have already begun), and the lower elevations will continue to dry out for several more weeks. Subsequently shallow landscape depressions where tactile foragers like ibis target prey species (crayfishes and small fishes), will have the longest periods of drying and potentially the highest THg concentrations.

4.2. Factors influencing egret and ibis chick mercury concentrations

Similar to adults, egret and ibis chicks differed considerably in the variables that influenced their Hg concentrations. Egret chick Hg blood levels were most influenced by date, whereas ibis chick Hg levels were most influenced by chick age, region, and water recession rate. The influence of date on egret Hg blood levels may be a factor of the age of chicks; as wading bird chicks get older they are often fed larger prey items (Ogden et al., 1976) that contain higher concentrations of Hg (Loftus 2000). Egret nests were highly synchronous across regions during our study; subsequently egret chick age and date were highly correlated. Date may not have been as important to ibis chick THg concentrations because ibis nesting was not as synchronous as that observed for egret chicks.

Variables that influenced ibis chick THg concentrations differed from factors that affected egret chicks. Whereas egret THg levels were only influenced by date, ibis THg levels were influenced by age, region, and water recession rates. Total Hg concentrations decreased with ibis




Fig. 5. Relationship between partial residuals of red blood cell Hg value ($\mu\text{g/g dw}$) and (A) age (d) of chicks at sampling and (B) water recession rate (cm/d) during the previous two weeks before sampling at adult foraging sites associated with colonies where white ibis chicks were sampled in the Florida Everglades, after accounting for the other predictor variables in the best models see Table 2). THg concentrations decreased by two-fold and increased by 162% across the range of ages and water recession rates, respectively (cumulative weights = 0.99 and 0.84, $\beta = -0.001 \pm 0.000$, $\beta = 0.248 \pm 0.201$, respectively).

chick age, suggesting that chicks likely depurated Hg into growing feathers or minimized their Hg burdens through mass dilution as they aged (Ackerman et al., 2011). Recent research on Forster's terns (*Sterna forsteri*), America avocets (*Recurvirostra americana*), and black-necked




Fig. 6. Conceptual model of the relationships between representative landscape environmental factors, foraging behavior, and related biogeochemical factors that may influence MeHg exposure in wading birds. Biogeochemical factors¹ were not assessed in modeling efforts.

stilts (*Himantopus mexicanus*) demonstrated that blood Hg concentrations followed a U-shaped pattern, Hg was highest just after chicks first hatched, decreased when they began to increase in size and depurate Hg into growing feathers, and then increased as both body growth and feather growth slowed (Ackerman et al., 2011). Previous research on Hg in ibis and egret chick feathers found a positive linear pattern in Hg concentrations (increasing with culmen length; a surrogate for age) in egrets but a negative linear pattern in ibises during the same two years of study in the Everglades (Herring et al., 2009). Similar to Ackerman et al. (2011), we found that ibis chick THg concentrations were elevated at a young age and then decreased considerably as they got older (3–12 days of age), were relatively stable in medium aged chicks (13–22 days of age), and actually decreased in older chicks (23–30 days of age). These differences in age-related Hg patterns may be associated with lower overall Hg exposure levels in Everglades' ibises relative to the avocets, stilts, and terns that were exposed to considerable higher levels of Hg in the San Francisco Bay (Ackerman et al., 2011). There is some evidence that ibis chick diet changes as chicks get older (switching to a diet with more crayfish). Dorn et al. (2011) found that at least during one year of our study (2007) in the Everglades, ibis chicks ate more fish earlier in the nestling period than at later stages of nesting, when crayfish were consumed more often than in the early nestling period. Lower Hg levels in crayfish relative to fish in the Everglades (Loftus, 2000) would then result in a decrease in Hg concentrations in old chicks prior to when their diet included more fish.

Similar to adult egrets, but not adult ibises, ibis chick THg concentrations were lower in Lox colonies than in WCA 3A colonies. These results are consistent with previous findings on feather Hg concentrations in ibis chicks in the Everglades during 2006 and 2007 (Herring et al., 2009). Further, as was the case in this study, Herring et al. (2009) found no differences in feather Hg concentrations in egret chicks between northern and southern regions. During the pre-breeding stage, adult ibises move around the landscape while searching for high quality sites (Beerens, 2008) and subsequently are not tied to specific regions as strongly as after they initiate nesting. Once they initiate nesting and begin provisioning chicks, the costs of foraging may restrict their movements to the localized region (e.g., Lox, WCA 3A) among which there are preexisting differences in Hg levels (Axelrad et al., 2007). It is unclear why egret chicks did not follow this pattern but overall egret chick THg patterns were consistent with egret chick feather Hg concentrations in Everglades in the same years (Herring et al., 2009).

Ibis chick THg concentrations were also influenced by water recession rates, increasing with positive recession rates (when water levels decrease). Positive water recession rates can result in decreased water depths over time, however in our study recession rates and water depths were not strongly correlated. Thus any effect recession rate had on ibis chicks is likely independent of the effect of decreased water depths resulting in prey becoming more abundant in shallow pools of water, and foraging adults consuming and subsequently feeding chicks potentially larger prey with higher Hg burdens. Previous research has shown that both recession rates and water depths determine when and where wading birds forage, however that relationship can change depending on levels of prey availability in the landscape (Beerens et al., 2011). We suspect that the relationship between ibis chick THg concentrations and recession rates is likely being driven by species-specific foraging patterns and foraging habitat selection processes (Beerens et al., 2011).

4.3. Egret and ibis Hg toxicity

To facilitate an understanding of egret and ibis adult and chick toxicity risk, we back-calculated whole blood THg values using linear model equation functions derived from a separate study and dataset on blood fraction THg concentrations in egret and ibis chicks ($R^2 = 0.63\text{--}0.85$; Herring, unpub data). The back-calculated geometric mean whole blood THg concentration ($\mu\text{g/g wet weight [ww]}$) was 0.33 in egret adults, 0.32 in egret chicks, 0.06 in ibis adults, and 0.03 in ibis

chicks. We assessed the potential for reproductive impairment in adults using toxicity values established in a recent experimental dosing study on ibis (Frederick and Jayasena, 2010). Based on our back-calculated values, both adult egrets and ibises on average were at low to no risk of impairment. Although toxicological threshold has not been developed for chicks, blood THg concentrations in egret and ibis chicks were lower than those of egret chicks in a dosing study where no effects were observed on plasma biomarkers (Hoffman et al., 2005). Further, at least in the case of egrets, chick blood THg concentrations at these sites have declined by over 60% since the mid 1990's (Sepulveda et al., 1999), suggesting that any risk for impairment of chicks has declined significantly during the last two decades. These findings are consistent with a reduction of external loading of Hg to the Everglades, however portions of the Everglades still have elevated Hg concentrations in fish, and wading birds may still be at risk of elevated Hg exposure (Rumbold et al., 2008).

5. Conclusions

Different landscape variables influenced THg concentrations across species and life stages in our study. Further, our results suggest differences among variables that influenced adults versus chicks, although we suggest caution when interpreting these results because the spatial and temporal differences in how the data were collected. For instance data in adults and chicks were collected from two separate time periods, resulting in potential differences in the hydrology and prey resources to which they were exposed to. Regardless, these results demonstrate for the first time that the hydrological conditions over temporal scales (in both the previous year and more recent [two weeks]) can play an important role in the subsequent Hg exposure of wading birds. The mechanisms associated with these two time scales may be different and are likely driven by the complex relationship associated with foraging ecology, landscape factors, and biogeochemical processes (Fig. 6).

Sites that regularly dry out provide an opportunity for methylation and mobilization of Hg from the surrounding landscape (Krabbenhoft and Fink, 2001; Osborne et al., 2011; Orem et al., 2011). Although drying of wetland sites in the Everglades is associated with the natural annual hydrologic cycle (Obeysekeru et al., 1999), the elevated atmospheric deposition of Hg onto the landscape (Osborne et al., 2011) has resulted in situations where wading birds may be continually exposed to Hg in this ecosystem dependent on the local hydrology and their foraging strategy. Further, in large portions of the Everglades, the numbers of days that sites are dry annually has increased due to the changes in hydrology associated with human alteration of the ecosystem (Davis et al., 1994; Bancroft et al., 2002; Lockwood et al., 2003). Although the exact relationship between the number of days a site was dry and the associated Hg in adult ibises is not clear, our results at least suggest that minimizing soil dry-out could aid in lowering MeHg production and wading bird Hg exposure (Krabbenhoft and Fink, 2001; Axelrad et al., 2007). Additionally, an improved understanding of how prey selection and associated Hg exposure may change with hydrologic conditions and across additional "searcher" and "exploiter" species may provide some clarity for this relationship.

Acknowledgments

Funding for research was provided by a grant to DEG and GH from the United States Fish and Wildlife Service and support from the U.S. Geological Survey. We appreciate the support and cooperation of the Arthur R. Marshall Loxahatchee National Wildlife Refuge staff. We thank our field research crews and fellow researchers that assisted in collection of field data and laboratory analyses: T. Anderson, N. Baker, T. Beck, B. Botson, K. Donner, H. Herring, N. Hill, A. Horton, B. Imdieke, B. Johnson, S. Lantz, and J. Pierce. M. Herzog kindly provided valuable advice and assistance on model selection and program R analyses. A. Jackson and M. Ricca provided valuable comments on a previous version of this manuscript. Florida Atlantic University Institutional Animal Care and Use Committee (Protocol A0534) approved the research techniques, and we

conducted the research under United States Fish and Wildlife Service Research Permit 23354 and Florida Fish and Wildlife Conservation Commission Scientific Research Permit WX04487. Any use of trade, product, or firm names is for descriptive purposes only and does not imply endorsement by the U.S. Government.

References

- Ackerman JT, Eagles-Smith CA, Herzog MP. Bird mercury concentrations change rapidly as chicks age: toxicological risk is highest at hatching and fledging. *Environ Sci Technol* 2011;45:5418–25.
- Axelrad DM, Atkeson TD, Lange T, Gilmour CC, Pollman CD, Orem WH, et al. Mercury monitoring, research and environmental assessment in South Florida. South Florida Environmental Report, Chapter 3B, South Florida Water Management District, West Palm Beach, FL; 2007. [57 pp.]
- Bancroft GT, Gawlik DE, Rutchev K. Distribution of wading birds relative to water vegetation and water depths in the northern Everglades of Florida, USA. *Waterbirds* 2002;25:265–77.
- Bearhop S, Ruxton GD, Furness RW. Dynamics of mercury in blood and feathers of great skuas. *Environ Toxicol Chem* 2000;19:1638–43.
- Beerens JM. Hierarchical resource selection and movements of two wading bird species with divergent foraging strategies in the Florida Everglades. [M.S. thesis] Boca Raton: Florida Atlantic University; 2008.
- Beerens JM, Gawlik DE, Herring G, Cook MI. Dynamic habitat selection by two wading bird species with divergent foraging strategies in a seasonally fluctuating wetland. *Auk* 2011;128:651–62.
- Burnham KP, Anderson DR. Model selection and inference: a practical information-theoretic approach. New York, New York: Springer-Verlag; 2002.
- Chen JW, Chang CC, Wang SL. Mercury concentration of whole blood and red blood cell in Taiwan residents. *Epidemiology* 2011;22:S289.
- Cleckner LB, Garrison PL, Hurley JP, Olson ML, Krabbenhoft DP. Trophic transfer of methyl mercury in the northern Florida Everglades. *Biogeochemistry* 1998;40:347–61.
- Crozier GE, Gawlik DE. The use of decoys in attracting wading birds. *J Field Ornithol* 2003;69:276–87.
- Custer TW, Custer CM, Johnson KM, Hoffman DJ. Mercury and other element exposure to tree swallows (*Tachycineta bicolor*) nesting on Lostwood National Wildlife Refuge, North Dakota. *Environ Pollut* 2008;155:217–26.
- Davis SM, Gunderson LH, Park WA, Richardson JR, Mattson JE. Landscape dimension, composition, and function in a changing Everglades ecosystem. In: Davis SM, Ogden JC, editors. Everglades: the ecosystem and its restoration. Delray Beach Florida: St. Lucie Press; 1994. p. 419–44.
- Day RD, Segars AL, Arendt MD, Lee AM, Peden-Adams MM. Relationship of blood mercury levels to health parameters in the loggerhead sea turtle (*Caretta caretta*). *Environ Health Perspect* 2007;115:1421–8.
- Dorn NJ, Cook MI, Herring G, Boyle R, Nelson J, Gawlik DE. Diet variation among white ibis (*Eudocimus albus*) chicks: prey composition depends on recent hydrologic conditions. *Ibis* 2011;153:323–35.
- Eagles-Smith CA, Ackerman JT, De La Cruz SE, Takekawa JY. Mercury bioaccumulation and risk to three waterbird foraging guilds is influenced by foraging ecology and breeding stage. *Environ Pollut* 2009;157:1993–2002.
- Fournier F, Karasov WH, Kenow KP, Meyer MW, Hines RK. The oral bioavailability and toxicokinetics of methylmercury in common loons (*Gavia immer*) chicks. *Comp Biochem Physiol A* 2002;133:703–14.
- Frederick PC, Collopy MW. Nesting success of Ciconiiform species in relation to water conditions in the Florida Everglades. *Auk* 1989;106:625–34.
- Frederick P, Jayasena N. Altered pairing behavior and reproductive success in white ibises exposed to environmentally relevant concentrations of methylmercury. *Proc Royal Soc Lon B* 2010;278:1851–7.
- Frederick PC, Spalding MG, Dusek K. Wading birds as bioindicators of mercury contamination in Florida: annual and geographic variation. *Environ Toxicol Chem* 2002;21:163–7.
- Gawlik DE. The effects of prey availability on the numerical response of wading birds. *Ecol Monog* 2002;72:329–46.
- Gilmour CC, Krabbenhoft D, Orem W, Aiken G. Appendix 2B-1: influence of drying and rewetting on mercury and sulfur cycling in Everglades and STA soils. 2004 Everglades Consolidated Report, South Florida Water Management District, West Palm Beach, FL; 2004.
- Gilmour CC, Krabbenhoft D, Orem W, Aiken G, Roden E. Appendix 3B-2. Status report on ACME studies on the control of mercury methylation and bioaccumulation in the Everglades. In: Redfield G, editor. 2007 South Florida Environmental Report, South Florida Water Management District, West Palm Beach FL; 2007.
- Gustin MS, Chavan PV, Dennett KE, Marchand EA, Donaldson S. Evaluation of wetland methyl mercury export as a function of experimental manipulations. *J Environ Qual* 2006;35:2352–9.
- Hall BD, Aiken GR, Krabbenhoft D, Marvin-DiPasquale M, Swarzenski CM. Wetlands as principal zones of methylmercury production in southern Louisiana and the Gulf of Mexico region. *Environ Pollut* 2008;154:124–34.
- Heath JA, Frederick PC. Trapping white ibis with rocket nets and mist nets in the Florida Everglades. *J Field Ornithol* 2003;69:276–87.
- Heath JA, Frederick PC. Relationships among mercury concentrations, hormones, and nesting effort of white ibises in the Florida Everglades. *Auk* 2005;122:255–67.
- Heath JA, Frederick PC, Kushlan JA, Bildstein KL. White ibis (*Eudocimus albus*), the birds of North America Online. In: Poole A, editor. Ithaca: Cornell Lab of ornithology; 2009. [Retrieved from the Birds of North America; <http://bna.birds.cornell.edu/bna/species/009>].

- Herring G, Cook MI, Gawlik DE, Call EM. Food availability is expressed through physiological stress indicators in nestling white ibis: a food supplementation experiment. *Func Ecol* 2011;25:682–90.
- Herring G, Gawlik DE. Differential physiological response to landscape prey availability by the great egret and white ibis. *J Wild Manage* 2012;77:58–67.
- Herring G, Gawlik DE, Beerens JM. Evaluating two new methods for capturing large wetland birds. *J Field Ornithol* 2008a;79:102–10.
- Herring G, Gawlik DE, Beerens JM. Predicting the sex of great egrets and white ibises. *Waterbirds* 2008b;31:306–11.
- Herring G, Gawlik DE, Cook MI, Beerens JM. Sensitivity of nesting great egrets (*Ardea alba*) and white ibises (*Eudocimus albus*) to reduced prey availability. *Auk* 2010;127:660–70.
- Herring G, Gawlik DE, Rumbold DA. Feather mercury concentrations and physiological condition of great egret and white ibis nestlings in the Florida Everglades. *Sci Total Environ* 2009;407:2641–9.
- Hoffman DJ, Spalding MG, Frederick PC. Subchronic effects of methylmercury on plasma and organ biochemistries in great egret nestlings. *Environ Toxicol Chem* 2005;24:3078–84.
- Jayasena N, Frederick PC, Larkin ILV. Endocrine disruption in white ibises (*Eudocimus albus*) caused by exposure to environmentally relevant levels of methylmercury. *Aquat Toxicol* 2011;105:321–7.
- Krabbenhoft DL, Fink LE. The effect of dry down and natural fires on mercury methylation in the Florida Everglades. In: Redfield G, editor. 2001 Everglades consolidated report. Report to the Florida Legislature West Palm Beach, FL: South Florida Water Management District; 2001. [A7-8-1–A7-8-14].
- Kushlan JA. Feeding ecology and prey selection in the white ibis. *Condor* 1979;81:376–89.
- Kushlan JA. Resource use strategies of wading birds. *Wilson Bull* 1981;93:145–63.
- Kushlan JA. Responses of wading birds to seasonally fluctuating water levels: strategies and their limits. *Colon Waterbirds* 1987;9:155–62.
- Lieske CL, Moses SK, Castellini JM, Kleka J, Hueffer K, O'Hara TM. Toxicokinetics of mercury in blood compartments and hair of fish-fed sled dogs. *Acta Vet Scand* 2011;53:66.
- Liu G, Cai Y, Phillip T, Kalla P, Scheidt D, Richards J, et al. Distribution of total and methylmercury in different ecosystem compartments in the Everglades implications for mercury bioaccumulation. *Environ Pollut* 2008;153:257–65.
- Liu Z, Volin JC, Owen VD, Pearlstine LG, Allen JR, Mazzotti FJ, et al. Validation and ecosystem applications of the EDEN water-surface model for the Florida Everglades. *Ecohydrol* 2009;2:182–94.
- Lockwood JL, Ross MS, Sah JP. Smoke on the water: the interplay of fire and water flow on Everglades Restoration. *Front Ecol Environ* 2003;1:462–8.
- Loftus WF. Accumulation and fate of mercury in an Everglade aquatic food web; doctoral thesis. Miami, FL: Florida International University; 2000.
- Marvin-DiPasquale MC, Agee JL, Bouse RM, Jaffe BE. Microbial cycling of mercury in contaminated pelagic and wetland sediments of San Pablo Bay, California. *Environ Geol* 2003;43:260–7.
- McCrimmon DA, Ogden JC, Bancroft GT. Great egret (*Ardea alba*). In: Poole A, Gill F, editors. *The birds of North America*, no. 570. Washington, D.C.: Academy of Natural Sciences, Philadelphia, and American Ornithologists' Union; 2011.
- Mercury Deposition Network. <http://nadp.sws.uiuc.edu/mdn/2012>.
- Molina CI, Gibon FM, Durbrey JL, Dominguez E, Guimarães JRD, Roulet M. Transfer of mercury and methylmercury along macroinvertebrate food chains in a floodplain lake of the Beni River, Bolivia Amazonia. *Sci Total Environ* 2010;408:3382–91.
- Myers GJ, Davidson PW. Prenatal methylmercury exposure and children: neurological development, and behavioral research. *Environ Health Perspect* 1998;106:841–7.
- Obeyssekera J, Browder J, Hornung L, Harwell MA. The natural South Florida system I: climate, geology, and hydrology. *Urban Ecosystems* 1999;3:223–44.
- Ogden JC. A comparison of wading bird nesting dynamics, 1931–1946 and 1974–1989 as an indication of changes in ecosystem conditions in the southern Everglades. In: Davis S, Ogden JC, editors. *Everglades: the Ecosystem and its Restoration* 1994. Delray Beach, FL, USA: St. Lucie Press; 1994. p. 533–70.
- Ogden JC, Davis SM, Brandt LA. Science strategy for a regional ecosystem monitoring and assessment program: the Florida Everglades example. In: Busch DE, Trexler JC, editors. *Monitoring Ecosystems: Interdisciplinary Approaches for Evaluating Ecoregional Initiatives*. Washington DC: Island Press; 2003. p. 135–63.
- Ogden JC, Kushlan JA, Tilmant JT. Prey selectivity by wood stork. *Condor* 1976;78:324–30.
- Orem W, Gilmour C, Axelrad D, Krabbenhoft D, Scheidt D, Kalla P, et al. Sulphur in the South Florida Ecosystem: distribution, sources, biogeochemistry, impacts, and management for restoration. *Crit Review Env Sci Tech* 2011;41:249–88.
- Osborne TZ, Newman S, Scheidt DJ, Kalla PI, Bruland GL, Cohen MJ, et al. Landscape patterns of significant soil nutrients and contaminants in the Greater Everglades Ecosystem: past, present, and future. *Crit Rev Environ Sci Tech* 2011;41:121–48.
- R Development Core Team. R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria3-900051-07-0; 2011. [URL <http://www.R-project.org/>].
- Rumbold DG, Niemczyk SL, Flink LE, Chandrasekhar T, Harkanson B, Kline KA. Mercury in eggs and feathers of great egrets (*Ardea alba*) from the Florida Everglades. *Arch Environ Contam Toxicol* 2001;41:501–7.
- Rumbold DG, Lange TR, Axelrad DM, Atkeson TD. Ecological risk of methylmercury in Everglades National Park, Florida, USA. *Ecotoxicology* 2008;17:632–41.
- Russell GJ, Bass Jr OL, Pimm SL. The effect of hydrological patterns and breeding-season flooding on the numbers and distribution of wading birds in Everglades National Park. *Anin Conserv* 2002;5:185–99.
- Sepulveda MS, Frederick PC, Spalding MG, Williams GE. Mercury contamination in free-ranging great egret nestlings (*Ardea alba*) from southern Florida. *Environ Toxicol Chem* 1999;18:985–92.
- Smith JP. Nesting season food habits of 4 species of herons and egrets at Lake Okeechobee, Florida. *Waterbirds* 1997;20:198–220.
- Sundlof SF, Spalding MG, Wentworth JD, Steible CK. Mercury in livers of wading birds (Ciconiiformes) in southern Florida. *Arch Environ Contam Toxicol* 1994;27:299–305.
- Thompson DR, Furness RW. Comparison of the levels of total and organic mercury in seabird feathers. *Mar Pollut Bull* 1989;20:577–9.
- U.S. Environmental Protection Agency. Method 7473, mercury in solids and solutions by thermal decomposition, amalgamation, and atomic absorption spectrophotometry, test methods for evaluating solid waste, physical/chemical methods SW 846, Update IVA. Washington, DC: U.S. Government Printing Office; 2000.
- US Geological Survey. The Everglades Depth Estimation Network (EDEN) for support of ecological and biological assessments. U. S. Department of the Interior Fact Sheet; 2006. p. 2006–3087.